



Available online at <http://scik.org>

J. Math. Comput. Sci. 5 (2015), No. 6, 822-825

ISSN: 1927-5307

A NOTE ON THE PAPER "COMMON FIXED POINT THEOREMS FOR THREE MAPS IN DISCONTINUOUS G_b -METRIC SPACES"

JING LIU, MEIMEI SONG*

College of Science, Tianjin University of Technology, Tianjin 300384, China

Copyright © 2015 Liu and Song. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. In this paper, fixed points three nonlinear operators are investigated. Common fixed point theorems are established in a complete G_b -metric space. The result presented in this paper improves the corresponding results in [1].

Keywords: G -metric space; b -metric space; Common fixed point.

2010 AMS Subject Classification: 47H10.

1. Main results

In [1], Roshan *et al.* obtained a common fixed point theorem in a complete G_b -metric space. After carefully reading the paper, the authors find that the proof of $d_{n+1} \leq d_n$ in Theorem 2.1 [1] turned out to be not comprehensive. They only proved $d_{3n+1} \leq d_{3n}$ and $d_{3n+2} \leq d_{3n+1}$ and declared that $d_{n+1} \leq d_n$, which has a skip.

Next, we give a new proof.

*Corresponding author

E-mail addresses: tjutliujing@hotmail.com (J. Liu), songmeimei@tjut.edu.cn (M. Song)

Received May 16, 2015

Theorem 1.1 Let (X, G) be a complete G_b -metric space. Let $A, B, C: X \rightarrow X$ be mappings which satisfy the following condition:

$$\psi(2s^4G(Ax, By, Cz)) \leq \psi(M(x, y, z)) - \varphi(M(x, y, z)) \quad (1.1)$$

for all $x, y, z \in X$, where $\psi, \varphi: [0, \infty) \rightarrow [0, \infty)$ are two mappings such that ψ is continuous nondecreasing, φ is a lower semi-continuous function with $\psi(t) = \varphi(t) = 0$ if and only if $t = 0$ and

$$M(x, y, z) = \max\{G(x, y, z), G(x, Ax, By), G(y, By, Cz), G(z, Cz, Ax)\}.$$

Then, either one of A, B , and C has a fixed point, or, the maps A, B and C have a unique common fixed point.

Proof. Choose $x_0 \in X$. Define the sequence $\{x_n\}$ as $x_{3n+1} = Ax_{3n}$, $x_{3n+2} = Bx_{3n+1}$ and $x_{3n+3} = Cx_{3n+2}$ for all $n = 0, 1, 2, \dots$. If $x_{3n} = x_{3n+1}$, then x_{3n} is a fixed point of A . If $x_{3n+1} = x_{3n+2}$, then x_{3n+1} is a fixed point of B . If $x_{3n+2} = x_{3n+3}$, then x_{3n+2} is a fixed point of C . Now, assume that $x_n \neq x_{n+1}$ for all n . Let $d_n = G(x_n, x_{n+1}, x_{n+2})$. we obtain from (1.1) that

$$\begin{aligned} \psi(d_{3n+1}) &\leq \psi(2s^4d_{3n+1}) = \psi(2s^4G(x_{3n+1}, x_{3n+2}, x_{3n+3})) \\ &= \psi(2s^4G(Ax_{3n}, Bx_{3n+1}, Cx_{3n+2})) \\ &\leq \psi(M(x_{3n}, x_{3n+1}, x_{3n+2})) - \varphi(M(x_{3n}, x_{3n+1}, x_{3n+2})), \end{aligned}$$

where

$$\begin{aligned} M(x_{3n}, x_{3n+1}, x_{3n+2}) &= \max\{G(x_{3n}, x_{3n+1}, x_{3n+2}), G(x_{3n}, Ax_{3n}, Bx_{3n+1}), \\ &\quad G(x_{3n+1}, Bx_{3n+1}, Cx_{3n+2}), G(x_{3n+2}, Cx_{3n+2}, Ax_{3n})\} \\ &= \max\{G(x_{3n}, x_{3n+1}, x_{3n+2}), G(x_{3n}, x_{3n+1}, x_{3n+2}), \\ &\quad G(x_{3n+1}, x_{3n+2}, x_{3n+3}), G(x_{3n+2}, x_{3n+3}, x_{3n+1})\} \\ &= \max\{d_{3n}, d_{3n}, d_{3n+1}, d_{3n+1}\} \\ &= \max\{d_{3n}, d_{3n+1}\}. \end{aligned}$$

We prove that $d_{3n+1} \leq d_{3n}$ for each $n \in \mathbb{N}$. If $d_{3n+1} > d_{3n}$ for some $n \in \mathbb{N}$, then we have $\psi(d_{3n+1}) \leq \psi(d_{3n+1}) - \varphi(d_{3n+1})$, which implies that $d_{3n+1} = 0$, a contradiction to $d_{3n+1} > 0$. Also, we have

$$\begin{aligned} \psi(d_{3n+2}) &\leq \psi(2s^4 d_{3n+2}) = \psi(2s^4 G(x_{3n+2}, x_{3n+3}, x_{3n+4})) \\ &= \psi(2s^4 G(Bx_{3n+1}, Cx_{3n+2}, Ax_{3n+3})) \\ &= \psi(2s^4 G(Ax_{3n+3}, Bx_{3n+1}, Cx_{3n+2})) \\ &\leq \psi(M(x_{3n+3}, x_{3n+1}, x_{3n+2})) - \varphi(M(x_{3n+3}, x_{3n+1}, x_{3n+2})), \end{aligned}$$

where

$$\begin{aligned} M(x_{3n+3}, x_{3n+1}, x_{3n+2}) &= \max\{G(x_{3n+3}, x_{3n+1}, x_{3n+2}), G(x_{3n+3}, Ax_{3n+3}, Bx_{3n+1}), \\ &\quad G(x_{3n+1}, Bx_{3n+1}, Cx_{3n+2}), G(x_{3n+2}, Cx_{3n+2}, Ax_{3n+3})\} \\ &= \max\{G(x_{3n+3}, x_{3n+1}, x_{3n+2}), G(x_{3n+3}, x_{3n+4}, x_{3n+2}), \\ &\quad G(x_{3n+1}, x_{3n+2}, x_{3n+3}), G(x_{3n+2}, x_{3n+3}, x_{3n+4})\} \\ &= \max\{d_{3n+1}, d_{3n+2}, d_{3n+1}, d_{3n+2}\} \\ &= \max\{d_{3n+1}, d_{3n+2}\}. \end{aligned}$$

Similarly, if $d_{3n+2} > d_{3n+1}$ for some $n \in \mathbb{N}$, then we have $\psi(d_{3n+2}) \leq \psi(d_{3n+2}) - \varphi(d_{3n+2})$, which implies that $d_{3n+2} = 0$, a contradiction to $d_{3n+2} > 0$. Also, we have

$$\begin{aligned} \psi(d_{3n+3}) &\leq \psi(2s^4 d_{3n+3}) = \psi(2s^4 G(x_{3n+3}, x_{3n+4}, x_{3n+5})) \\ &= \psi(2s^4 G(Cx_{3n+2}, Ax_{3n+3}, Bx_{3n+4})) \\ &= \psi(2s^4 G(Ax_{3n+3}, Bx_{3n+4}, Cx_{3n+2})) \\ &\leq \psi(M(x_{3n+3}, x_{3n+4}, x_{3n+2})) - \varphi(M(x_{3n+3}, x_{3n+4}, x_{3n+2})), \end{aligned}$$

where

$$\begin{aligned}
 M(x_{3n+3}, x_{3n+4}, x_{3n+2}) &= \max\{G(x_{3n+3}, x_{3n+4}, x_{3n+2}), G(x_{3n+3}, Ax_{3n+3}, Bx_{3n+4}), \\
 &\quad G(x_{3n+4}, Bx_{3n+4}, Cx_{3n+2}), G(x_{3n+2}, Cx_{3n+2}, Ax_{3n+3})\} \\
 &= \max\{G(x_{3n+3}, x_{3n+4}, x_{3n+2}), G(x_{3n+3}, x_{3n+4}, x_{3n+5}), \\
 &\quad G(x_{3n+4}, x_{3n+5}, x_{3n+3}), G(x_{3n+2}, x_{3n+3}, x_{3n+4})\} \\
 &= \max\{d_{3n+2}, d_{3n+3}, d_{3n+3}, d_{3n+2}\} \\
 &= \max\{d_{3n+2}, d_{3n+3}\}.
 \end{aligned}$$

Similarly, if $d_{3n+3} > d_{3n+2}$ for some $n \in \mathbb{N}$, then we have $\psi(d_{3n+3}) \leq \psi(d_{3n+3}) - \varphi(d_{3n+3})$, which implies that $d_{3n+3} = 0$, a contradiction to $d_{3n+3} > 0$. Hence, we have $0 < d_{n+1} \leq d_n$ for each $n \in \mathbb{N}$. The rest proof process is the same with which was given in [1]. We, therefore, omit the proof.

Conflict of Interests

The authors declare that there is no conflict of interests.

REFERENCES

- [1] J.R. Roshan *et al.*, Common fixed point theorems for three maps in discontinuous G_b -metric spaces, *Acta Math. Sci.* 34 (2014), 1643-1654.