



Available online at <http://scik.org>

Adv. Fixed Point Theory, 8 (2018), No. 2, 205-220

<https://doi.org/10.28919/afpt/3637>

ISSN: 1927-6303

ON THE TOPOLOGY OF n -VALUED MAPS

ROBERT F. BROWN^{1,*}, DACIBERG LIMA GONÇALVES²

¹Department of Mathematics, University of California, Los Angeles, CA 90095-1555, USA

²Departamento de Matemática - IME, Universidade de São Paulo, CEP 05508-090 - São Paulo - SP - Brazil

Copyright © 2018 Brown and Gonçalves. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. This paper presents an exposition of the topological foundations of the theory of n -valued maps. By means of proofs that exploit particular features of n -valued functions, as distinct from more general classes of multivalued functions, we establish, among other properties, the equivalence of several definitions of continuity. The exposition includes an exploration of the role of configuration spaces in the study of n -valued maps. As a consequence of this point of view, we extend the classical Splitting Lemma, that is central to the fixed point theory of n -valued maps, to a characterization theorem that leads to a new type of construction of non-split n -valued maps.

Keywords: n -valued map; splitting lemma; continuity; graph of a function; configuration space; braid group.

2010 AMS Subject Classification: 54C60, 55M20, 57M10, 20F36.

*Corresponding author

E-mail address: rfb@math.ucla.edu

Received January 2, 2018

1. Introduction

An n -valued map $\phi : X \multimap Y$ is a continuous multivalued function that associates to each $x \in X$ an unordered subset of exactly n points of Y . The fixed point theory of n -valued maps $\phi : X \multimap X$ has been a topic of considerable interest in recent years and there is much research activity at the present time [3] - [12], [15], [16]. The purpose of this paper is to present the topological foundations of this subject.

In Section 2 we assume as few hypotheses on the topological spaces X and Y as necessary in order to obtain the results. For a general multivalued function, continuity is defined as the satisfaction of two independent conditions called upper and lower semi-continuity. The section begins by proving that, in the case of n -valued functions, lower semi-continuity implies upper semi-continuity (but not the converse) and thus this single condition is sufficient for the definition of an n -valued map. We use this result to prove that, in the setting of n -valued functions, the general definition of continuity for multivalued functions is equivalent to a classical definition based on convergence of sequences.

The *graph* $\Gamma(\phi)$ of a multivalued function $\phi : X \multimap Y$ is the set

$$\Gamma(\phi) = \{(x, y) \in X \times Y : y \in \phi(x)\},$$

topologized as a subspace of $X \times Y$. The graph plays a significant role in the general theory of multivalued functions, see for instance [17]. If ϕ is an n -valued map, then its graph has a very useful property: the projection $p_X : \Gamma(\phi) \rightarrow X$ defined by $p_X(x, y) = x$ is a finite covering space. This covering space has played a key role in the construction of n -valued maps.

In Section 3 we characterize the splitting of an n -valued map that extends the classical Splitting Lemma. An n -valued map $\phi : X \multimap Y$ is *split* if there are single-valued maps $f_1, \dots, f_n : X \rightarrow Y$ such that $\phi(x) = \{f_1(x), \dots, f_n(x)\}$ for all $x \in X$. The Splitting Lemma states that if X is simply connected and locally path connected, then every n -valued map $\phi : X \multimap Y$ is split. The Splitting Lemma was first proved by Banach and Mazur in [1] to obtain conditions that imply

that a local homeomorphism is a global homeomorphism.¹ The Nielsen fixed point theory of an n -valued map $\phi: X \multimap X$ presented by Schirmer in [23] - [25] depends on a version of the Splitting Lemma which she obtained from a more general result concerning multivalued functions on simply connected compact metric spaces due to O'Neill [21]. Another consequence of the Splitting Lemma, in [6], is that an n -valued map $\phi: X \multimap Y$ of finite simplicial complexes induces a chain map of their simplicial chain complexes. It follows that the integer Lefschetz number of an n -valued self-map of a finite simplicial complex is well-defined and the Lefschetz Fixed Point Theorem can be generalized to such maps.

The configuration space $D_n(Y)$ of a space Y is the set of all unordered sets of n points of Y . The set of n -valued functions from a space X to a space Y is in one-to-one correspondence with the set of single-valued functions from X to $D_n(Y)$ where, to a given n -valued function $\phi: X \multimap Y$ we associate the single-valued function $\Phi: X \rightarrow D_n(Y)$ defined by $\Phi(x) = \{\phi(x)\}$. We prove in Section 3 that, under suitable hypotheses, ϕ is an n -valued map if and only if Φ is continuous with respect to a natural topology on $D_n(Y)$.²

Thus for $\phi: X \multimap Y$ an n -valued map, the map Φ induces a homomorphism $\Phi_\#: \pi_1(X) \rightarrow \pi_1(D_n(Y))$. We prove that this homomorphism characterizes the splitting of n -valued maps. If X is simply connected, we obtain the Splitting Lemma as a consequence of this result. Moreover, we demonstrate that hypotheses on X more general than simple connectedness can be sufficient to imply that all n -valued maps $\phi: X \multimap Y$ are split.

The group $\pi_1(D_n(Y))$ may be identified with the braid group of n strands on Y [13]. This identification facilitates the construction in Section 3 of non-split n -valued self-maps $\phi: X \multimap X$

¹The proof in [1] is for n -valued maps of metric spaces. However, the authors state that the same result can be generalized to topological spaces.

²The configuration space viewpoint was introduced into the fixed point theory of n -valued maps in [15] and [16]. However, configuration spaces have been a well-established tool in the fixed point and coincidence theory of single-valued maps, see for instance [14] where the configuration space of ordered pairs is presented as the complement of the diagonal of the Cartesian square of a space.

that are of interest in the fixed point theory of such maps, complementing the constructions in [12] and [8].

Finally, in Section 4 we add the hypothesis that the range Y of an n -valued function $\phi : X \multimap Y$ is a metric space. Therefore, for $x, x' \in X$, the Hausdorff distance between $\phi(x)$ and $\phi(x')$ as subsets of Y is well-defined. We can use the Hausdorff distance to define a form of continuity for ϕ and we prove that, in this setting, it is equivalent to the other definitions of continuity that we have presented. As a tool for this proof, we use the concept of the gap of an n -valued map that was introduced by Schirmer in [23] and we establish its properties.

Most of the results in this paper are not new and are actually special cases of established facts. The n -valued maps belong to a class of multivalued functions called weighted maps, and these have been extensively studied [22]. Our goal is to furnish a person interested in the topic of n -valued maps with arguments for their basic topological properties that are as simple and elementary as possible, without any dependence on a more general theory.

However, although the primary purpose of this paper is the exposition of known facts about the topology of n -valued maps, in the course of its preparation some interesting features of such maps appeared that, it seems, have not been noted previously. We call the reader's attention to the property of n -valued maps presented as Proposition 2.1, that lower semi-continuity implies upper semi-continuity. This property not only simplifies the proofs of some of the subsequent results, it is quite special to n -valued maps, as an example in Section 2 demonstrates. Moreover, while the Splitting Lemma of Corollary 3.1 has been a basic tool since the initiation of the fixed point theory of n -valued maps by Schirmer, the Characterization Theorem 3.1 is a considerable extension of this classical result and, as demonstrated in Section 3, it has several significant consequences.

The authors thank Ofelia Alas for clarifying for them the relationship of the continuity of an n -valued function and the continuity of a mapping to the corresponding configuration space.

2. Functions of Hausdorff spaces

Throughout this section, we assume only that the spaces are Hausdorff.

A multivalued function $\phi : X \multimap Y$ is *lower semi-continuous* if V an open subset of Y implies that the set $\{x \in X : \phi(x) \cap V \neq \emptyset\}$ is open in X and *upper semi-continuous* if V open in Y implies that the set $\{x \in X : \phi(x) \subseteq V\}$ is open in X . Since we will need terminology that distinguishes several definitions of continuity, we will call ϕ *multicontinuous* if it is both lower and upper semi-continuous.³

Let $\phi : X \multimap Y$ be a lower semi-continuous function and $x^{(0)} \in X$ with $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$. Let V_1, \dots, V_n be disjoint open subsets of Y such that $y_j^{(0)} \in V_j$. Since ϕ is lower semi-continuous, for each j there is an open subset U_j of X containing $x^{(0)}$ such that if $x \in U_j$, then $\phi(x) \cap V_j \neq \emptyset$ and we define $U(x^{(0)}, \{V_j\}) = \bigcap_{j=1}^n U_j$. If $x \in U(x^{(0)}, \{V_j\})$, then $\phi(x) \subseteq \bigcup_{j=1}^n V_j$ and we can number the points of $\phi(x) = \{y_1, \dots, y_n\}$ so that $y_j \in V_j$.

Proposition 2.1. *Let $\phi : X \multimap Y$ be an n -valued function. If ϕ is lower semi-continuous it is also upper semi-continuous and therefore multicontinuous, that is, an n -valued map.*

Proof. Let $x^{(0)} \in X$ and let V be an open subset of Y containing $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$. Let V_1, \dots, V_n be disjoint open subsets of V such that $y_j^{(0)} \in V_j$ for each j . If $x \in U(x^{(0)}, \{V_j\})$, then $\phi(x) \subseteq \bigcup_{j=1}^n V_j \subseteq V$ so $\{x \in X : \phi(x) \subseteq V\}$ is open in X and we have proved that ϕ is upper semi-continuous. \square

To demonstrate that an upper semi-continuous n -valued function need not be continuous, define a 2-valued function $\phi : \mathbb{R} \multimap \mathbb{R}$ by $\phi(x) = \{x, x/2\}$ if $x \neq 0$ and $\phi(0) = \{0, 1\}$. If $V \subseteq \mathbb{R}$ is an open set containing $\phi(0)$, then there exists $\varepsilon > 0$ such that $(-\varepsilon, \varepsilon) \subseteq V$ and if $|x| < \varepsilon$, then $\phi(x) \subset (-\varepsilon, \varepsilon) \subseteq V$. Since ϕ is continuous at $x \neq 0$ and thus upper semi-continuous, we

³In Wikipedia, these concepts are called upper and lower *hemi-continuous* and the words upper and lower semi-continuous are reserved for certain generalizations of continuity for single-valued functions which, together, imply continuity in the usual sense. However, in both [2] and [17] the word semi-continuous is used as it is here.

conclude that ϕ is an upper semi-continuous function. However,

$$\{x \in \mathbb{R} : \phi(x) \cap (1/2, 3/2) \neq \emptyset\} = (1/2, 3) \cup \{0\}$$

which is not open. Therefore ϕ is not lower semi-continuous and thus not continuous.

The property of n -valued maps presented in Proposition 2.1 fails immediately beyond the class of n -valued functions. For instance, define an “almost 2-valued function” $\phi: [0, 1] \multimap [0, 1]$ by $\phi(x) = \{0, 1\}$ if $x \neq 0$ and $\phi(0) = \{0\}$. Then ϕ is lower semi-continuous because if U_0 and U_1 are neighborhoods of 0 and 1 respectively, then $\{x: \phi(x) \cap U_0 \neq \emptyset\} = [0, 1]$ and $\{x: \phi(x) \cap U_1 \neq \emptyset\} = (0, 1]$. But $\{x: \phi(x) \subseteq U_0\} = \{0\}$ so ϕ is not upper semi-continuous.

The function $\phi: X \multimap Y$ is *locally split* if, given $x^{(0)} \in X$ there is a neighborhood U of $x^{(0)}$ and maps $g_1, \dots, g_n: U \rightarrow Y$ such that $\phi(x) = \{g_1(x), \dots, g_n(x)\}$ for all $x \in U$.

Proposition 2.2. *An n -valued map $\phi: X \multimap Y$ is locally split.*

Proof. Let $x^{(0)} \in X$ with $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$ and V_1, \dots, V_n disjoint open subsets of Y with $y_j^{(0)} \in V_j$. For $x \in U = U(x^{(0)}, \{V_j\})$ with $\phi(x) = \{y_1, \dots, y_n\}$, we number the y_j such that $y_j \in V_j$. For $j = 1, \dots, n$, define $g_j: U \rightarrow Y$ by $g_j(x) = y_j$, then $\phi(x) = \{g_1(x), \dots, g_n(x)\}$. To prove that the g_j are continuous, let $x_* \in U$ and let W be an open subset of Y that contains $g_j(x_*)$. Let $Z = \{x \in X : \phi(x) \cap (W \cap V_j) \neq \emptyset\}$ which contains x_* and is open because ϕ is lower semi-continuous. Then $U \cap Z$ is an open subset of X containing x_* such that if $x \in U \cap Z$, then $g_j(x) = \phi(x) \cap (W \cap V_j) \subseteq W$. \square

It follows easily from Proposition 2.2 that

Proposition 2.3. *If $\phi: X \multimap Y$ is an n -valued map, then the map $p_X: \Gamma(\phi) \rightarrow X$ defined by $p_X(x, y) = x$ is a covering space.*

Thus an n -valued map determines a finite covering space. Conversely, a finite covering space determines an n -valued map, as follows.

Proposition 2.4. *If $p: \tilde{X} \rightarrow X$ is a covering space of degree n , then $p^{-1}: X \multimap \tilde{X}$ is an n -valued map.*

Proof. Let U be an open subset of \tilde{X} and let $x_0 \in X$ such that $p^{-1}(x_0) \cap U \neq \emptyset$. Because $p: \tilde{X} \rightarrow X$ is a covering space, then p is an open map so $p(U)$ is an open subset of X containing x_0 . Since $\phi(x) \cap U \neq \emptyset$ for all $x \in p(U)$, we conclude that p^{-1} is a lower semi-continuous function and therefore, by Proposition 2.1, an n -valued map. \square

The n -valued map p^{-1} is used in [8] to construct n -valued self-maps of \tilde{X} and in [12] to construct n -valued self-maps of X .

In [1], Banach and Mazur called an n -valued function $\phi: X \multimap Y$ with $\phi(x) = \{y_1, \dots, y_n\}$ continuous if, given a sequence $x^{(k)}$ in X such that $\lim_{k \rightarrow \infty} x^{(k)} = x^{(0)}$ with $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$, the sets $\phi(x^{(k)})$ can be ordered as $y_1^{(k)}, \dots, y_n^{(k)}$ so that $\lim_{k \rightarrow \infty} y_i^{(k)} = y_i^{(0)}$ for all $i = 1, \dots, n$. We will call an n -valued function that satisfies this definition *sequentially continuous*.

As another application of Proposition 2.1 we have

Proposition 2.5. *If an n -valued function $\phi: X \multimap Y$ is sequentially continuous and X is a first-countable space, then ϕ is multicontinuous.*

Proof. It follows from Proposition 2.1 that ϕ is multicontinuous if and only if it is lower semi-continuous. Suppose ϕ is not lower semi-continuous, then there exists $x^{(0)} \in X$ and an open subset V of Y such that $\phi(x^{(0)}) \cap V \neq \emptyset$ but, for any open subset U of X containing $x^{(0)}$, there a point $x \in U$ such that $\phi(x) \cap V = \emptyset$. Since X is first countable, there is a basis $\mathcal{U} = \{U_k\}_{k=1}^{\infty}$ for the topology of X at $x^{(0)}$. For each k , there is a point $x^{(k)} \in U_k$ such that $\phi(x^{(k)}) \cap V = \emptyset$. Let $y_j^{(0)} \in \phi(x^{(0)})$ such that $y_j^{(0)} \in V$. Since \mathcal{U} is a basis for the topology of X at $x^{(0)}$, then $\lim_{k \rightarrow \infty} x^{(k)} = x^{(0)}$, but $\phi(x^{(k)}) = \{y_1^{(k)}, \dots, y_n^{(k)}\}$ cannot be ordered so that $\lim_{k \rightarrow \infty} y_j^{(k)} = y_j^{(0)}$, and therefore ϕ is not sequentially continuous. \square

The converse of the previous result holds for all Hausdorff spaces:

Proposition 2.6. *If an n -valued function $\phi : X \multimap Y$ is multicontinuous, then it is sequentially continuous.*

Proof. Let $x^{(0)} \in X$ such that $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$. Let $(x^{(k)})$ be a sequence in X such that $\lim_{k \rightarrow \infty} x^{(k)} = x^{(0)}$. Suppose $V_j \subseteq Y$ are open sets such that $y_j^{(0)} \in V_j$. We may assume, without loss of generality, that the subsets V_j are disjoint. Since ϕ is lower semi-continuous, we observed at the beginning of this section that therefore there is an open set $U(x^{(0)}, \{V_j\})$ in X containing $x^{(0)}$ such that if $x \in U(x^{(0)}, \{V_j\})$, then $\phi(x) = \{y_1, \dots, y_j\}$ can be ordered so that $y_j \in V_j$. Since $\lim_{k \rightarrow \infty} x^{(k)} = x^{(0)}$, there exists N such that if $k > N$, then $x^{(k)} \in U(x^{(0)}, \{V_j\})$. Therefore, if $k > N$ then $y_j^{(k)} \in V_j$, so $\lim_{k \rightarrow \infty} y_j^{(k)} = y_j^{(0)}$ and we conclude that ϕ is sequentially continuous. \square

3. Configuration spaces and the characterization of splitting

As in the previous section, we assume only that all spaces are Hausdorff unless other conditions are stated.

Given a space Y and a positive integer n , we define $F_n(Y)$, the *ordered configuration space* of Y , by

$$F_n(Y) = \{(y_1, \dots, y_n) \in Y^n = Y \times \dots \times Y : y_i \neq y_j \text{ for } i \neq j\}.$$

The symmetric group S_n acts freely on $F_n(Y)$ and the quotient space $D_n(Y) = F_n(Y)/S_n$ is the *unordered configuration space* of Y . Topologize $F_n(Y)$ as a subset of the product space Y^n and then give $D_n(Y)$ the quotient topology so that, for the quotient map $q : F_n(Y) \rightarrow D_n(Y)$, a subset W of $D_n(Y)$ is open if and only if $q^{-1}(W)$ is open in $F_n(Y)$.

It is observed in [15] that the set of n -valued functions from a space X to a space Y is in one-to-one correspondence with the set of single-valued functions from X to $D_n(Y)$. Specifically, if $\phi : X \multimap Y$ is an n -valued function, then the corresponding function $\Phi : X \rightarrow D_n(Y)$ is defined by $\Phi(x) = \{\phi(x)\}$.

Proposition 3.1. *If $\phi: X \multimap Y$ is an n -valued map, then the corresponding function $\Phi: X \rightarrow D_n(Y)$ is continuous.*

Proof. Let $x^{(0)} \in X$ then, by Proposition 2.2, there is a neighborhood U of $x^{(0)}$ and single-valued maps $g_1, \dots, g_n: U \rightarrow Y$ such that $\phi(x) = \{g_1(x), \dots, g_n(x)\}$ for all $x \in U$. Therefore, the restriction of $\Phi: X \rightarrow D_n(Y)$ to U is the composition of the function $\hat{\phi}: U \rightarrow F_n(x)$ defined by $\hat{\phi}(x) = (g_1(x), \dots, g_n(x))$ and the projection $q: F_n(Y) \rightarrow D_n(Y)$, both of which are continuous. Therefore, the function Φ is continuous. \square

Proposition 3.1 allows us to replace an n -valued map of spaces by a single-valued map, the range of which is an unordered configuration space.

We will illustrate the configuration space concept by describing the ordered and unordered configuration spaces of the circle S^1 . In addition to the relative simplicity of this setting, it is natural to focus on it because the n -valued self-maps of the circle may be viewed as the motivating example for this branch of fixed point theory. The reciprocal of the square-root function on the complex numbers of norm one was introduced in [24]: it is a non-split 2-valued self-map of the circle such that every 2-valued map of the circle homotopic to it has at least three fixed points.⁴

Our presentation is based on a paper of Westerland [27] and comments of Dylan Thurston [26]. We begin with the ordered configuration space $F_n(S^1)$. This space is the disjoint union of $(n-1)!$ sets determined by the $n!/n = (n-1)!$ orderings of n points z_1, \dots, z_n of S^1 . For a given ordering, choose the value of z_1 as the S^1 coordinate, then the successive differences of the polar coordinates between the adjacent points in the cyclic ordering of the points determines $n-1$ positive real numbers whose sum is less than one and so they define an open $n-1$ -simplex. Thus each component of $F_n(S^1)$ is homeomorphic to the product of the circle and an open $n-1$ -simplex and consequently it is the homotopy type of the circle.

⁴The fixed point theory of n -valued maps of the circle was subsequently developed in [5].

The quotient map $q: F_n(S^1) \rightarrow D_n(S^1)$ is a covering space of order $n!$ and the restriction of q to each component of $F_n(S^1)$ is of order n . Thus $D_n(S^1)$ is a $K(\pi, 1)$ because $F_n(S^1)$ is. To determine the fundamental group π of $D_n(S^1)$, we first note that that group must be torsion free since $D_n(S^1)$ is a finite-dimensional manifold and it admits the cyclic group \mathbb{Z} as a subgroup of finite index. Since π is a virtually cyclic group that is torsion free, the classification of virtually cyclic groups implies that $\pi = \mathbb{Z}$. The homomorphism of cyclic groups $q_\#: \pi_1(F_n(S^1)) \rightarrow \pi_1(D_n(S^1))$ induced by q is multiplication by n .

The converse of Proposition 3.1 also holds, as follows.

Proposition 3.2. *Let $\phi: X \multimap Y$ be an n -valued function. If X is locally path-connected and semilocally simply connected and $\Phi: X \rightarrow D_n(Y)$ is continuous, then ϕ is multicontinuous, that is, an n -valued map.⁵*

Proof. Let $x^{(0)} \in X$ be an arbitrary point then, since X is locally path-connected and semilocally simply connected, there is a path-connected neighborhood U of $x^{(0)}$ such that $i_\#: \pi_1(U) \rightarrow \pi_1(X)$, the homomorphism induced by the inclusion, is trivial. Therefore the homomorphism $(\Phi|U)_\#: \pi_1(U) \rightarrow \pi_1(D_n(Y))$ induced by the restriction of Φ to U is trivial because it factors through $i_\#$. By [18], Proposition 1.33, the map $\Phi|U$ admits a lifting to $F_n(Y)$ and we can view it as consisting of maps $g_1, \dots, g_n: U \rightarrow Y$ such that $g_i(x) \neq g_j(x)$ for $i \neq j$. Therefore the restriction to U of the corresponding n -valued function $\phi: X \multimap Y$ is split by the maps g_j . Let $x^{(0)} \in U$ and let V be an open subset of Y that contains some $g_j(x^{(0)})$. Since g_j is continuous, there exists an open neighborhood W of $x^{(0)}$ in U such that $g_j(W) \subseteq V$. Therefore $\phi(x) \cap V \neq \emptyset$ for all $x \in W$ and we have proved that ϕ is lower semi-continuous which, by Proposition 2.1, implies that ϕ is an n -valued map. \square

⁵By means of a more elaborate proof, based on Corollary 9.3 of [20], it can be demonstrated that this proposition does not require the hypotheses that the domain is locally path-connected and semilocally simply connected. However, such a proof is beyond the scope of the present paper.

By Propositions 3.1 and 3.2, if the domain satisfies appropriate hypotheses, we obtain another characterization of continuity for n -valued functions, as follows:

Corollary 3.1. *If X is locally path-connected and semilocally simply connected, an n -valued function $\phi: X \multimap Y$ is multicontinuous if and only if the corresponding function $\Phi: X \rightarrow D_n(Y)$ is continuous.*

By [13] the fundamental group of the configuration space $D_n(Y)$ is $B_n(Y)$, the braid group of n strands on Y , and the fundamental group of $F_n(Y)$ is $P_n(Y)$, the subgroup of $B_n(Y)$ of pure braids. Since if $\phi: X \multimap Y$ is an n -valued map, then the corresponding function $\Phi: X \rightarrow D_n(Y)$ is continuous, so it induces a homomorphism $\Phi_{\#}: \pi_1(X) \rightarrow B_n(Y)$ (compare [15]). We use that homomorphism to obtain the following generalization of the Splitting Lemma.

Theorem 3.1. (*Splitting Characterization Theorem*) *Let $\phi: X \multimap Y$ be an n -valued map, where X is connected and locally path-connected. Then ϕ is split if and only if the image of $\Phi_{\#}: \pi_1(X) \rightarrow B_n(Y)$ is contained in the image of the homomorphism $q_{\#}: P_n(Y) \rightarrow B_n(Y)$ induced by the projection $q: F_n(Y) \rightarrow D_n(Y)$. In particular, if X is simply connected, then all n -valued maps $\phi: X \multimap Y$ are split.⁶*

Proof. Suppose that ϕ is split. This implies that there exist maps $g_1, \dots, g_n: X \rightarrow Y$ such that $\phi(x) = \{g_1(x), \dots, g_n(x)\}$ and we may define a map $\hat{\phi}: X \rightarrow F_n(Y)$ by $\hat{\phi}(x) = (g_1(x), \dots, g_n(x))$. Then the corresponding map $\Phi: X \rightarrow D_n(Y)$ is the composition of $\hat{\phi}$ with the projection $q: F_n(Y) \rightarrow D_n(Y)$ and it follows that the image of the induced homomorphism $\Phi_{\#}: \pi_1(X) \rightarrow B_n(Y)$ is contained in the image of $q_{\#}: P_n(Y) \rightarrow B_n(Y)$.

Suppose that the image of the homomorphism $\Phi_{\#}: \pi_1(X) \rightarrow B_n(Y)$ is contained in the image of $q_{\#}: P_n(Y) \rightarrow B_n(Y)$, the homomorphism induced by the projection $q: F_n(Y) \rightarrow D_n(Y)$. By

⁶This form of the Splitting Lemma corresponds to that of Banach and Mazur [1] for Hausdorff spaces. The proof of Schirmer [23], based on a result of O'Neill [21], does not require that the domain be locally path-connected, but it only applies when both domain and range are compact spaces.

[18], Proposition 1.33 there is a lifting $\hat{\phi}: X \rightarrow F_n(Y)$ of Φ . The coordinate maps of $\hat{\phi}$ split ϕ and the result follows. \square

Not only must n -valued maps with domain a simply connected space be split but, more generally, if X and Y are spaces such that there are no nontrivial homomorphism from $\pi_1(X)$ to $B_n(Y)$, then Theorem 3.1 implies that every n -valued map from X to Y must be split. For example, let G be a finite group, let X_G be a 2-complex such that $\pi_1(X_G) = G$ and let Y be a surface that is not the projective plane then, since $B_n(Y)$ is a torsion free group, all n -valued maps $\phi: X_G \rightarrow Y$ must be split.

Viewing a braid β in $B_n(Y)$ as an n -valued map of the unit interval to Y where $\beta(1)$ is a permutation of $\beta(0)$, define $\rho: B_n(Y) \rightarrow S_n$ by sending β to the corresponding permutation. Since the braid group sequence

$$1 \rightarrow P_n(Y) \xrightarrow{q\#} B_n(Y) \xrightarrow{\rho} S_n \rightarrow 1$$

is exact ([19], page 16), then the Splitting Characterization Theorem implies that if $\phi: X \rightarrow Y$ induces $\Phi\#: \pi_1(X) \rightarrow B_n(Y)$ such that the composition $\rho\Phi\#: \pi_1(X) \rightarrow S_n$ is nontrivial, then ϕ does not split.

From the point of view of fixed point theory, the study of split n -valued self-maps of a finite polyhedron can be reduced to the single-valued setting because Schirmer proved in [24] that if $\phi = \{f_1, \dots, f_n\}: X \rightarrow X$, then the Nielsen number of ϕ is calculated from that of the f_i by the formula $N(\phi) = \sum_{i=1}^n N(f_i)$. On the other hand, examples from [8] demonstrate that $N(\phi)$ can exhibit a variety of behaviors in the non-split case. This motivates the use of our Splitting Characterization Theorem to construct non-split n -valued self-maps $\phi: X \rightarrow X$ which we will do by defining ϕ so that the image of $\Phi\#$ does not lie in the subgroup $q\#(P_n(Y)) \subseteq B_n(Y)$ of pure braids on Y . In order to make the construction, the only requirement is that the first Betti number of X be non-zero. Thus the integer cohomology group $H^1(X)$ is of rank at least one. Identifying $H^1(X)$ with the Buschinsky group of homotopy classes of maps $[X, S^1]$, that group is nontrivial so we may choose a map $a: X \rightarrow S^1$ which induces an epimorphism $a\#: \pi_1(X) \rightarrow \pi_1(S^1)$.

Let $\beta \in B_n(X)$ be a braid on X such that $\rho(\beta) \in S_n$ is not the identity permutation. Since $B_n(Y) = \pi_1(D_n(Y))$, the homotopy class $[b]$ corresponding to the braid β is represented by a map $b: S^1 \rightarrow D_n(X)$ which may be viewed as an n -valued map $b: S^1 \multimap X$. Then $\phi = b \circ a: X \multimap X$ is a non-split n -valued map.

4. Maps to a metric range

Let $\phi: X \multimap Y$ be an n -valued function and write $\phi(x) = \{y_1, \dots, y_n\}$ for $x \in X$. For Y a metric space with metric d_Y , define $\gamma_\phi: X \rightarrow \mathbf{R}$ by

$$\gamma_\phi(x) = \min_{1 \leq i \neq j \leq n} d_Y(y_i, y_j).$$

In [23], the *gap* $\gamma(\phi)$ of ϕ is defined as

$$\gamma(\phi) = \inf\{\gamma_\phi(x) : x \in X\}.$$

The following useful fact is stated in [23] without a proof. We take this opportunity to present its brief proof.

Proposition 4.1. *Let Y be a metric space with metric d_Y and let $\phi: X \multimap Y$ be a multicontinuous n -valued function, then the function $\gamma_\phi: X \rightarrow \mathbf{R}$ is continuous. Therefore, if X is compact, then the gap $\gamma(\phi) > 0$ for all ϕ .*

Proof. Let $x_0 \in X$, then by Proposition 2.2 there is a neighborhood U of x_0 and maps $g_1, \dots, g_n: U \rightarrow Y$ such that $\phi(x) = \{g_1(x), \dots, g_n(x)\}$ for all $x \in U$. Then for $x \in U$, the gap is defined by

$$\gamma_\phi(x) = \min_{1 \leq i \neq j \leq n} \{d_Y(g_i(x), g_j(x))\}.$$

Since the g_i are continuous, γ_ϕ is continuous on U and therefore it is a continuous function. \square

The gap, and in particular the fact that it is positive on compact spaces, is used in [23] to obtain a simplicial approximation theorem for n -valued maps and, subsequently, in [6] to define a simplicial chain map induced by the n -valued map. It appears also in the proof of the

uniqueness of splitting in [9] and in the reduction of the computation of the Nielsen number of an n -valued map to a coincidence number for single-valued maps in [10].

Let $\phi : X \multimap Y$ be an n -valued function where Y is a metric space with metric d_Y . For $x, x' \in X$, the *Hausdorff distance*, when specialized to that between $\phi(x) = \{y_1, \dots, y_n\}$ and $\phi(x') = \{y'_1, \dots, y'_n\}$, denoted $d_H(\phi(x), \phi(x'))$, is the greater of

$$\max_{1 \leq j \leq n} \min_{1 \leq i \leq n} d_Y(y_j, y'_i)$$

and

$$\max_{1 \leq i \leq n} \min_{1 \leq j \leq n} d_Y(y_j, y'_i).$$

We define $\phi : X \multimap Y$ to be *Hausdorff continuous* at $x_0 \in X$ if, given $\varepsilon > 0$, there exists an open subset U of X containing x_0 such that if $x \in U$, then $d_H(\phi(x_0), \phi(x)) < \varepsilon$. It is *Hausdorff continuous on X* if it is Hausdorff continuous at every point of X .

Proposition 4.2. *An n -valued function $\phi : X \multimap Y$, where Y is a metric space, is multicontinuous if and only if it is Hausdorff continuous on X .*

Proof. We first prove that if $\phi : X \multimap Y$ is Hausdorff continuous, then it is multicontinuous. By Proposition 2.1, we need only prove that ϕ is lower semi-continuous. Let $x^{(0)} \in X$ such that $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$ and let $V \subseteq Y$ be an open set such that $\phi(x^{(0)}) \cap V \neq \emptyset$; we assume $y_1^{(0)} \in V$. Choose $\varepsilon > 0$ so that $B(y_1^{(0)}, \varepsilon) \subseteq V$ then, since ϕ is Hausdorff continuous, there is an open subset U of X containing $x^{(0)}$ such that $x \in U$ implies that $d_H(\phi(x), \phi(x^{(0)})) < \varepsilon$. Therefore, for $\phi(x) = \{y_1, \dots, y_j\}$ we know that

$$\max_{1 \leq i \leq n} \min_{1 \leq j \leq n} d_Y(y_j, y_i^{(0)}) < \varepsilon.$$

and, in particular, that

$$\min_{1 \leq j \leq n} d_Y(y_j, y_1^{(0)}) < \varepsilon$$

so some $y_j \in B(y_1^{(0)}, \varepsilon)$. Therefore, $x \in U$ implies that $\phi(x) \cap V \neq \emptyset$ and we have demonstrated that the set of $x \in X$ with this property is open.

Now suppose that $\phi: X \multimap Y$ is multicontinuous, that $x^{(0)} \in X$, and we are given $\varepsilon > 0$. We may assume that $\varepsilon < \gamma_\phi(x^{(0)})/2$. Let $\phi(x^{(0)}) = \{y_1^{(0)}, \dots, y_n^{(0)}\}$ and define disjoint open sets $V_j = B(y_j^{(0)}, \varepsilon)$. We will show that if x is in the open subset $U(x^{(0)}, \{V_j\})$ of X defined in Section 2, then $d_H(\phi(x), \phi(x^{(0)})) < \varepsilon$ so ϕ is Hausdorff continuous at x_0 . Let $\phi(x) = \{y_1, \dots, y_n\}$. If $i \neq j$, then

$$\begin{aligned} \gamma_\phi(x^{(0)}) &\leq d_Y(y_i^{(0)}, y_j^{(0)}) \\ &\leq d_Y(y_i^{(0)}, y_j) + d_Y(y_j, y_j^{(0)}) \\ &< d_Y(y_i^{(0)}, y_j) + \gamma_\phi(x^{(0)})/2 \end{aligned}$$

so

$$d_Y(y_i^{(0)}, y_j) > \gamma_\phi(x^{(0)})/2.$$

Therefore, since $d_Y(y_i^{(0)}, y_i) < \gamma_\phi(x^{(0)})/2$ for all i , we conclude that

$$d_H(\phi(x), \phi(x^{(0)})) = \max_{1 \leq i \leq n} d_Y(y_i^{(0)}, y_i) < \varepsilon.$$

□

Conflict of Interests

The authors declare that there is no conflict of interests.

REFERENCES

- [1] Banach, S. and Mazur, S., *Über mehrdeutige stetige Abbildungen*, Studia Math., **5** (1934), 174 - 178.
- [2] Berge, C., *Topological Spaces*, Olivier & Boyd, 1963.
- [3] Better, J., *Equivariant Nielsen fixed point theory for n -valued maps*, Topology Appl., **157** (2010), 1804 - 1814.
- [4] Better, J., *A Wecken theorem for n -valued maps*, Topology Appl., **159** (2012), 3707 - 3715.
- [5] Brown, R., *Fixed points of n -valued multimaps of the circle*, Bull. Polish Acad. Sci. Math., **54** (2006), 153 - 162.
- [6] Brown, R., *The Lefschetz number of an n -valued multimap*, JP Journal of Fixed Point Theory Appl., **2** (2007), 53 - 60.

- [7] Brown, R., *Nielsen numbers of n -valued fiber maps*, J. Fixed Point Theory Appl., **4** (2008), 183 - 201.
- [8] Brown, R., *Construction of multiply fixed n -valued maps*, Topology Appl., **196** (2015), 249 - 259.
- [9] Brown, R. and Kolahi, K., *Nielsen coincidence, fixed point and root theories of n -valued maps*, J. Fixed Point Theory Appl., **14** (2013), 309 - 324.
- [10] Brown, R. and Lin, J., *Coincidences of projections and linear n -valued maps of tori*, Topology Appl., **157** (2010), 1990 - 1998.
- [11] Brown, R. and Nan, J., *Stabilizers of fixed point classes and Nielsen numbers of n -valued maps*, Bull. Belgian Math. Soc., **24** (2017), 523-535.
- [12] Crabb, M., *Lefschetz indices for n -valued maps*, J. Fixed Point Theory Appl., **17** (2015), 153 - 186.
- [13] Fox, R. and Neuwirth, L., *The braid groups*, Math. Scand., **10** (1962), 119 - 126.
- [14] Gonçalves, D., *Braid groups and Wecken pairs*, Contemp. Math., **72** (1988), 89 - 97.
- [15] Gonçalves, D. and Guaschi, J., *Fixed points of multimaps, the fixed point property and the case of surfaces - a braid approach*, Indag. Math., **29** (2018), 91 - 124.
- [16] Gonçalves, D. and Guaschi, J., *Fixed points of n -valued maps on surfaces and the Wecken property - configuration space approach*, Science China Mathematics, **60** (2017), 1561 - 1574.
- [17] Gorniewicz, L., *Topological Fixed Point Theory of Multivalued Mappings*, Kluwer Academic Publishers, 1999.
- [18] Hatcher, A., *Algebraic Topology*, Cambridge U. Press, 2002.
- [19] Hansen, V., *Braids and coverings*, London Math. Soc., 1989.
- [20] Michael, E., *Topologies for spaces of subsets*, Trans. Amer. Math. Soc., **17** (1951), 152 - 182.
- [21] O'Neill, B., *Induced homology homomorphisms for set-valued maps*, Pacific J. Math., **7** (1957), 1179 - 1184.
- [22] Pejsachowicz, J. and Skiba, R., *Fixed point theory of multivalued weighted maps*, Handbook of Topological Fixed Point Theory, Springer, 217 - 263 (2005).
- [23] Schirmer, H., *Fix-finite approximations of n -valued multifunctions*, Fund. Math., **121** (1984), 73 - 80.
- [24] Schirmer, H., *An index and Nielsen number for n -valued multifunctions*, Fund. Math., **124** (1984), 207 - 219.
- [25] Schirmer, H., *A minimum theorem for n -valued multifunctions*, Fund. Math., **126** (1985), 83 - 92.
- [26] Thurston, D., <https://mathoverflow.net/questions/206546/unordered-configuration-space-of-pointed-space>.
- [27] Westerland, C., *Configuration spaces in topology and geometry*, Gaz. Australian Math. Soc., **38** (2011), 279 - 283.