



Available online at <http://scik.org>

J. Math. Comput. Sci. 8 (2018), No. 3, 304-317

<https://doi.org/10.28919/jmcs/3003>

ISSN: 1927-5307

## GENERALIZED $\sigma$ -CONVERGENT DIFFERENCE SEQUENCE SPACES DEFINED BY ORLICZ FUNCTION

KHALID EBADULLAH

College of Science and Theoretical Studies, Saudi Electronic University, Kingdom of Saudi Arabia

Copyright © 2018 Khalid Ebadullah. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Abstract.** In this paper, we introduce the sequence space  $V_{\sigma}(M, p, r, \Delta^u)$ , where  $u \in N$ ,  $M$  is an Orlicz function,  $p = (p_m)$  is any sequence of strictly positive real numbers and  $r \geq 0$  and study some of the properties and inclusion relations that arise on the said space.

**Keywords:** invariant mean; paranorm; orlicz function and difference sequences.

**2010 AMS Subject Classification:** 40F05, 40C05, 46A45.

### 1. Introduction

Let  $N$ ,  $R$  and  $C$  be the sets of all natural, real and complex numbers respectively.

We write

$$\omega = \{x = (x_k) : x_k \in R \text{ or } C\},$$

the space of all real or complex sequences.

Let  $\ell_{\infty}$ ,  $c$  and  $c_0$  denote the Banach spaces of bounded, convergent and null sequences respectively.

---

E-mail address: [khalidebadullah@gmail.com](mailto:khalidebadullah@gmail.com)

Received October 14, 2016

The following subspaces of  $\omega$  were first introduced and discussed by Maddox [13-14].

$$\ell(p) = \{x \in \omega : \sum_k |x_k|^{p_k} < \infty\},$$

$$\ell_\infty(p) = \{x \in \omega : \sup_k |x_k|^{p_k} < \infty\},$$

$$c(p) = \{x \in \omega : \lim_k |x_k - l|^{p_k} = 0, \text{ for some } l \in C\},$$

$$c_0(p) = \{x \in \omega : \lim_k |x_k|^{p_k} = 0\},$$

where  $p = (p_k)$  is a sequence of strictly positive real numbers.

The concept of paranorm is closely related to linear metric spaces. It is a generalization of that of absolute value. (see [14])

Let  $X$  be a linear space. A function  $g : X \rightarrow R$  is called paranorm, if for all  $x, y, z \in X$ ,

$$(P1) \quad g(x) = 0 \text{ if } x = \theta,$$

$$(P2) \quad g(-x) = g(x),$$

$$(P3) \quad g(x + y) \leq g(x) + g(y),$$

(P4) If  $(\lambda_n)$  is a sequence of scalars with  $\lambda_n \rightarrow \lambda$  ( $n \rightarrow \infty$ ) and  $x_n, a \in X$  with  $x_n \rightarrow a$  ( $n \rightarrow \infty$ ), in the sense that  $g(x_n - a) \rightarrow 0$  ( $n \rightarrow \infty$ ), in the sense that  $g(\lambda_n x_n - \lambda a) \rightarrow 0$  ( $n \rightarrow \infty$ ).

An Orlicz function is a function  $M : [0, \infty) \rightarrow [0, \infty)$ , which is continuous, non-decreasing and convex with  $M(0) = 0$ ,  $M(x) > 0$  for  $x > 0$  and  $M(x) \rightarrow \infty$  as  $x \rightarrow \infty$ .

Lindenstrauss and Tzafriri [11] used the idea of Orlicz functions to construct the sequence space

$$\ell_M = \{x \in \omega : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty, \text{ for some } \rho > 0\}$$

The space  $\ell_M$  is a Banach space with the norm

$$\|x\| = \inf\{\rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1\}$$

The space  $\ell_M$  is closely related to the space  $\ell_p$  which is an Orlicz sequence space with  $M(x) = x^p$  for  $1 \leq p < \infty$ .

An Orlicz function  $M$  is said to satisfy  $\Delta_2$  condition for all values of  $x$  if there exists a constant  $K > 0$  such that  $M(Lx) \leq KLM(x)$  for all values of  $L > 1$ .

A sequence space  $E$  is said to be solid or normal if  $(x_k) \in E$  implies  $(\alpha_k x_k) \in E$  for all sequence of scalars  $(\alpha_k)$  with  $|\alpha_k| < 1$  for all  $k \in N$ .

For Orlicz function and related results see([1],[2],[12], [17-21]).

Let  $\sigma$  be an injection on the set of positive integers  $N$  into itself having no finite orbits and  $T$  be the operator defined on  $\ell_\infty$  by  $T(x_k) = (x_{\sigma(k)})$ .

A positive linear functional  $\Phi$ , with  $\|\Phi\| = 1$ , is called a  $\sigma$ -mean or an invariant mean if  $\Phi(x) = \Phi(Tx)$  for all  $x \in \ell_\infty$ .

A sequence  $x$  is said to be  $\sigma$ -convergent, denoted by  $x \in V_\sigma$ , if  $\Phi(x)$  takes the same value, called  $\sigma - \lim x$ , for all  $\sigma$ -means  $\Phi$ . We have

$$V_\sigma = \{x = (x_k) : \sum_{m=1}^{\infty} t_{m,n}(x) = L \text{ uniformly in } n, L = \sigma - \lim x\},$$

where for  $m \geq 0, n > 0$ .

$$t_{m,n}(x) = \frac{x_k + x_{\sigma(k)} + \dots + x_{\sigma^m(k)}}{m+1}, \text{ and } t_{-1,n} = 0.$$

where  $\sigma^m(k)$  denotes the  $m^{\text{th}}$  iterate of  $\sigma$  at  $n$ . In particular, if  $\sigma$  is the translation, a  $\sigma$ -mean is often called a Banach limit and  $V_\sigma$  reduces to  $f$ , the set of almost convergent sequences.

Subsequently the spaces of invariant mean and Orlicz function have been studied by various authors. See([1],[12],[17-18],[21]).

The idea of Difference sequence sets

$$X_\Delta = \{x = (x_k) \in \omega : \Delta x = (x_k - x_{k+1}) \in X\},$$

where  $X = \ell_\infty, c$  or  $c_0$  was introduced by Kizmaz [10].

Kizmaz [10] defined the sequence spaces,

$$\ell_\infty(\Delta) = \{x = (x_k) \in \omega : (\Delta x_k) \in \ell_\infty\},$$

$$c(\Delta) = \{x = (x_k) \in \omega : (\Delta x_k) \in c\},$$

$$c_0(\Delta) = \{x = (x_k) \in \omega : (\Delta x_k) \in c_0\},$$

where  $\Delta x = (x_k - x_{k+1})$ . These are Banach spaces with the norm

$$\|x\|_\Delta = |x_1| + \|\Delta x\|_\infty.$$

After then Mikael [15] defined the sequence spaces :

$$\ell_\infty(\Delta^2) = \{x = (x_k) \in \omega : (\Delta^2 x_k) \in \ell_\infty\},$$

$$c(\Delta^2) = \{x = (x_k) \in \omega : (\Delta^2 x_k) \in c\},$$

$$c_0(\Delta^2) = \{x = (x_k) \in \omega : (\Delta^2 x_k) \in c_0\},$$

and showed that these are Banach spaces with norm

$$\|x\|_\Delta = |x_1| + |x_2| + \|\Delta^2 x\|_\infty.$$

After then Colak and Mikael[16] defined the sequence spaces

$$\ell_\infty(\Delta^m) = \{x = (x_k) \in \omega : (\Delta^m x_k) \in \ell_\infty\},$$

$$c(\Delta^m) = \{x = (x_k) \in \omega : (\Delta^m x_k) \in c\},$$

$$c_0(\Delta^m) = \{x = (x_k) \in \omega : (\Delta^m x_k) \in c_0\},$$

where  $m \in N$ ,

$$\Delta^0 x = (x_k),$$

$$\Delta x = (x_k - x_{k+1}),$$

$$\Delta^m x = (\Delta^{m-1} x_k - \Delta^{m-1} x_{k+1}),$$

and so that

$$\Delta^m x_k = \sum_{i=0}^m (-1)^i \begin{bmatrix} m \\ i \end{bmatrix} x_{k+i}.$$

and showed that these are Banach spaces with the norm

$$\|x\|_{\Delta} = \sum_{i=1}^m |x_i| + \|\Delta^m x\|_{\infty}.$$

For difference sequences see([3-9],[10],[15],[16]).

Recently Ebadullah[6] introduced and studied the sequence space

$$V_{\sigma}(M, p, r) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}.$$

Where  $M$  is an Orlicz function,  $p = (p_m)$  is any sequence of strictly positive real numbers and  $r \geq 0$ .

After then Ebadullah[7] introduced the sequence space

$$V_{\sigma}(M, p, r, \Delta) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}.$$

and discussed the following sequence spaces ;

For  $M(x) = x$  we get

$$V_{\sigma}(p, r, \Delta) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} |t_{m,n}(\Delta x)|^{p_m} < \infty \text{ uniformly in } n\}$$

For  $p_m = 1$ , for all  $m$ , we get

$$V_{\sigma}(M, r, \Delta) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta x)|}{\rho})] < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $r = 0$  we get

$$V_{\sigma}(M, p, \Delta) = \{x = (x_k) : \sum_{m=1}^{\infty} [M(\frac{|t_{m,n}(\Delta x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $M(x) = x$  and  $r=0$  we get

$$V_{\sigma}(p, \Delta) = \{x = (x_k) : \sum_{m=1}^{\infty} |t_{m,n}(\Delta x)|^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $p_k = 1$ , for all  $m$  and  $r=0$ , we get

$$V_{\sigma}(M, \Delta) = \{x = (x_k) : \sum_{m=1}^{\infty} [M(\frac{|t_{m,n}(\Delta x)|}{\rho})] < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $M(x) = x$ ,  $p_m = 1$ , for all  $m$  and  $r=0$ , we get

$$V_{\sigma}(\Delta x) = \{x = (x_k) : \sum_{m=1}^{\infty} |t_{m,n}(\Delta x)| < \infty \text{ uniformly in } n\}.$$

Later on Ebadullah[8] introduce the sequence space

$$V_{\sigma}(M, p, r, \Delta^2) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^2 x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}.$$

and studied the following sequence spaces ;

For  $M(x) = x$  we get

$$V_{\sigma}(p, r, \Delta^2) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} |t_{m,n}(\Delta^2 x)|^{p_m} < \infty \text{ uniformly in } n\}$$

For  $p_m = 1$ , for all  $m$ , we get

$$V_{\sigma}(M, r, \Delta^2) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^2 x)|}{\rho})] < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $r = 0$  we get

$$V_{\sigma}(M, p, \Delta^2) = \{x = (x_k) : \sum_{m=1}^{\infty} [M(\frac{|t_{m,n}(\Delta^2 x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $M(x) = x$  and  $r=0$  we get

$$V_{\sigma}(p, \Delta^2) = \{x = (x_k) : \sum_{m=1}^{\infty} |t_{m,n}(\Delta^2 x)|^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $p_k = 1$ , for all  $m$  and  $r=0$ , we get

$$V_{\sigma}(M, \Delta^2) = \{x = (x_k) : \sum_{m=1}^{\infty} [M(\frac{|t_{m,n}(\Delta^2 x)|}{\rho})] < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $M(x) = x$ ,  $p_m = 1$ , for all  $m$  and  $r=0$ , we get

$$V_{\sigma}(\Delta^2 x) = \{x = (x_k) : \sum_{m=1}^{\infty} |t_{m,n}(\Delta^2 x)| < \infty \text{ uniformly in } n\}.$$

## 2. Main results

In this article we introduce the sequence space

$$V_{\sigma}(M, p, r, \Delta^u) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}.$$

Where  $M$  is an Orlicz function,  $u \in N$ ,  $p = (p_m)$  is any sequence of strictly positive real numbers and  $r \geq 0$ .

Now we define the sequence spaces as follows;

For  $M(x) = x$  we get

$$V_{\sigma}(p, r, \Delta^u) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} |t_{m,n}(\Delta^u x)|^{p_m} < \infty \text{ uniformly in } n\}$$

For  $p_m = 1$ , for all  $m$ , we get

$$V_{\sigma}(M, r, \Delta^u) = \{x = (x_k) : \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})] < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $r = 0$  we get

$$V_{\sigma}(M, p, \Delta^u) = \{x = (x_k) : \sum_{m=1}^{\infty} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $M(x) = x$  and  $r=0$  we get

$$V_{\sigma}(p, \Delta^u) = \{x = (x_k) : \sum_{m=1}^{\infty} |t_{m,n}(\Delta^u x)|^{p_m} < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $p_k = 1$ , for all  $m$  and  $r=0$ , we get

$$V_{\sigma}(M, \Delta^u) = \{x = (x_k) : \sum_{m=1}^{\infty} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})] < \infty \text{ uniformly in } n, \rho > 0\}$$

For  $M(x) = x$ ,  $p_m = 1$ , for all  $m$  and  $r=0$ , we get

$$V_{\sigma}(\Delta^u x) = \{x = (x_k) : \sum_{m=1}^{\infty} |t_{m,n}(\Delta^u x)| < \infty \text{ uniformly in } n\}.$$

**Theorem 2.1.** The sequence space  $V_{\sigma}(M, p, r, \Delta^u)$  is a linear space over the field  $C$  of complex numbers.

**Proof.** Let  $x, y \in V_{\sigma}(M, p, r, \Delta^u)$  and  $\alpha, \beta \in C$  then there exists positive numbers  $\rho_1$  and  $\rho_2$  such that

$$\sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho_1})]^{p_m} < \infty,$$

and

$$\sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u y)|}{\rho_2})]^{p_m} < \infty$$

uniformly in  $n$ .

Define  $\rho_3 = \max(2|\alpha|\rho_1, 2|\beta|\rho_2)$ .

Since  $M$  is non decreasing and convex we have

$$\begin{aligned} & \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|\alpha t_{m,n}(\Delta^u x) + \beta t_{m,n}(\Delta^u y)|}{\rho_3})]^{p_m} \\ & \leq \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|\alpha t_{m,n}(\Delta^u x)|}{\rho_3} + \frac{|\beta t_{m,n}(\Delta^u y)|}{\rho_3})]^{p_m} \\ & \leq \sum_{m=1}^{\infty} \frac{1}{m^r} \frac{1}{2} [M(\frac{t_{m,n}(\Delta^u x)}{\rho_1}) + M(\frac{t_{m,n}(\Delta^u y)}{\rho_2})] < \infty \end{aligned}$$



uniformly in  $n$ .

This proves that  $V_{\sigma}(M, p, r, \Delta^u)$  is a linear space over the field  $\mathbb{C}$  of complex numbers.

**Theorem 2.2.** For any Orlicz function  $M$  and a bounded sequence  $p = (p_m)$  of strictly positive real numbers,  $V_{\sigma}(M, p, r, \Delta^u)$  is a paranormed space with

$$g(x) = \inf_{n \geq 1} \left\{ \rho^{\frac{pn}{H}} : \left( \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} \right)^{\frac{1}{H}} \leq 1, \text{ uniformly in } n \right\}$$

where  $H = \max(1, \sup p_m)$ .

**Proof.** It is clear that  $g(\Delta^u x) = g(-\Delta^u x)$ .

Since  $M(0) = 0$ , we get

$$\inf \left\{ \rho^{\frac{pn}{H}} \right\} = 0, \text{ for } x = 0$$

Now for  $\alpha = \beta = 1$ , we get

$$g(\Delta^u x + \Delta^u y) \leq g(\Delta^u x) + g(\Delta^u y).$$

For the continuity of scalar multiplication let  $l \neq 0$  be any complex number. Then by the definition we have

$$g(l\Delta^u x) = \inf_{n \geq 1} \left\{ \rho^{\frac{pn}{H}} : \left( \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(l\Delta^u x)|}{\rho})]^{p_m} \right)^{\frac{1}{H}} \leq 1, \text{ uniformly in } n \right\}$$

$$g(l\Delta^u x) = \inf_{n \geq 1} \left\{ (|l|s)^{\frac{pn}{H}} : \left( \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(l\Delta^u x)|}{(|l|s)})]^{p_m} \right)^{\frac{1}{H}} \leq 1, \text{ uniformly in } n \right\}$$

where  $s = \frac{\rho}{|l|}$ .

Since  $|l|^{p_m} \leq \max(1, |l|^H)$ , we have

$$g(l\Delta^u x) \leq \max(1, |l|^H) \inf_{n \geq 1} \left\{ s^{\frac{pn}{H}} : \left( \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{(|l|s)})]^{p_m} \right)^{\frac{1}{H}} \leq 1, \text{ uniformly in } n \right\}$$

$$g(\Delta^u l x) \leq \max(1, |l|^H) g(\Delta^u x)$$

Therefore  $g(\Delta^u x)$  converges to zero when  $g(\Delta^u x)$  converges to zero in  $V_\sigma(M, p, r, \Delta^u)$ .

Now let  $x$  be fixed element in  $V_\sigma(M, p, r, \Delta^u)$ . There exists  $\rho > 0$  such that

$$g(\Delta^u x) = \inf_{n \geq 1} \{ \rho^{\frac{pn}{H}} : (\sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m})^{\frac{1}{H}} \leq 1, \text{ uniformly in } n \}$$

Now

$$g(l\Delta^u x) = \inf_{n \geq 1} \{ \rho^{\frac{pn}{H}} : (\sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(l\Delta^u x)|}{\rho})]^{p_m})^{\frac{1}{H}} \leq 1, \text{ uniformly in } n \} \rightarrow 0 \text{ as } l \rightarrow 0.$$

This completes the proof.

**Theorem 2.3.** Suppose that  $0 < p_m < t_m < \infty$  for each  $m \in N$  and  $r > 0$ . Then

- (a)  $V_\sigma(M, p, \Delta^u) \subseteq V_\sigma(M, t, \Delta^u)$ .
- (b)  $V_\sigma(M, \Delta^u) \subseteq V_\sigma(M, r, \Delta^u)$

**Proof.**(a) Suppose that  $x \in V_\sigma(M, p, \Delta^u)$ .

This implies that  $[M(\frac{|t_{i,n}(\Delta^u x)|}{\rho})]^{p_m} \leq 1$

for sufficiently large value of  $i$ , say  $i \geq m_0$  for some fixed  $m_0 \in N$ .

Since  $M$  is non decreasing, we have

$$\sum_{m=m_0}^{\infty} [M(\frac{|t_{i,n}(\Delta^u x)|}{\rho})]^{t_m} \leq \sum_{m=m_0}^{\infty} [M(\frac{|t_{i,n}(\Delta^u x)|}{\rho})]^{p_m} < \infty.$$

Hence  $x \in V_\sigma(M, t, \Delta^u)$ .

(b) The proof is trivial.

**Corollary 2.4.**  $0 < p_m \leq 1$  for each  $m$ , then  $V_\sigma(M, p, \Delta^u) \subseteq V_\sigma(M, \Delta^u)$

If  $p_m \geq 1$  for all  $m$ , then  $V_\sigma(M, \Delta^u) \subseteq V_\sigma(M, p, \Delta^u)$ .

**Theorem 2.5.** The sequence space  $V_\sigma(M, p, r, \Delta^u)$  is solid.

**Proof.** Let  $x \in V_\sigma(M, p, r, \Delta^u)$ . This implies that

$$\sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} < \infty.$$

Let  $\alpha_m$  be a sequence of scalars such that  $|\alpha_m| \leq 1$  for all  $m \in N$ . Then the result follows from the following inequality.

$$\sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|\alpha_m t_{i,n}(\Delta^u x)|}{\rho})]^{p_m} \leq \sum_{m=1}^{\infty} \frac{1}{m^r} [M(\frac{|t_{i,n}(\Delta^u x)|}{\rho})]^{p_m} < \infty.$$

Hence  $\alpha x \in V_\sigma(M, p, r, \Delta^u)$  for all sequence of scalars  $(\alpha_m)$  with  $|\alpha_m| \leq 1$  for all  $m \in N$  whenever  $x \in V_\sigma(M, p, r, \Delta^u)$ .

**Corollary 2.6.** The sequence space  $V_\sigma(M, p, r, \Delta^u)$  is monotone.

**Theorem 2.7.** Let  $M_1, M_2$  be Orlicz function satisfying  $\Delta_2$  condition and  $r, r_1, r_2 \geq 0$ . Then we have

- (a) If  $r > 1$  then  $V_\sigma(M_1, p, r, \Delta^u) \subseteq V_\sigma(M \circ M_1, p, r, \Delta^u)$ ,
- (b)  $V_\sigma(M_1, p, r, \Delta^u) \cap V_\sigma(M_2, p, r, \Delta^u) \subseteq V_\sigma(M_1 + M_2, p, r, \Delta^u)$ ,
- (c) If  $r_1 \leq r_2$  then  $V_\sigma(M, p, r_1, \Delta^u) \subseteq V_\sigma(M, p, r_2, \Delta^u)$ .

**Proof.** (a) Since  $M$  is continuous at 0 from right, for  $\varepsilon > 0$  there exists  $0 < \delta < 1$  such that  $0 \leq c \leq \delta$  implies  $M(c) < \varepsilon$ .

If we define

$$I_1 = \{m \in N : M_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho}) \leq \delta \text{ for some } \rho > 0\},$$

$$I_2 = \{m \in N : M_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho}) > \delta \text{ for some } \rho > 0\},$$

when

$$M_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho}) > \delta$$

we get

$$M(M_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})) \leq \{\frac{2M(1)}{\delta}\}M_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})$$

Hence for  $x \in V_\sigma(M_1, p, r, \Delta^u)$  and  $r > 1$

$$\begin{aligned} \sum_{m=1}^{\infty} \frac{1}{m^r} [MOM_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} &= \sum_{m \in I_1} \frac{1}{m^r} [MOM_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} + \sum_{m \in I_2} \frac{1}{m^r} [MOM_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m}. \\ \sum_{m=1}^{\infty} \frac{1}{m^r} [MOM_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} &\leq \max(\varepsilon^h, \varepsilon^H) \sum_{m=1}^{\infty} \frac{1}{m^r} + \max(\{\frac{2M_1}{\delta}\}^h, \{\frac{2M_1}{\delta}\}^H) \end{aligned}$$

$$\text{where } 0 < h = \inf p_m \leq p_m \leq H = \sup p_m < \infty$$

(b)The proof follows from the following inequality

$$\frac{1}{m^r} [(M_1 + M_2)(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} \leq C \frac{1}{m^r} [M_1(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m} + C \frac{1}{m^r} [M_2(\frac{|t_{m,n}(\Delta^u x)|}{\rho})]^{p_m}$$

(c) The proof is straightforward.

**Corollary 2.8.** Let  $M$  be an Orlicz function satisfying  $\Delta_2$  condition. Then we have

- (a) If  $r > 1$  then  $V_\sigma(p, r, \Delta^u) \subseteq V_\sigma(M, p, r, \Delta^u)$ ,
- (b)  $V_\sigma(M, p, \Delta^u) \subseteq V_\sigma(M, p, r, \Delta^u)$ ,
- (c)  $V_\sigma(p, \Delta^u) \subseteq V_\sigma(p, r, \Delta^u)$ ,
- (d)  $V_\sigma(M, \Delta^u) \subseteq V_\sigma(M, r, \Delta^u)$ .

**Proof.** The proof is straightforward.

### Conflict of Interests

The authors declare that there is no conflict of interests.

### REFERENCES

- [1] Ahmad, Z.U., Mursaleen, M. An application of Banach limits. *Proc.Amer. Math. soc.* 103 (1983), 244-246.
- [2] Chen, S.T. Geometry of Orlicz spaces. *Dissertation Math. The Institute of Mathematics, Polish Academy of Sciences.*(1996)
- [3] Ebadullah, K. On certain class of sequence spaces of invariant mean defined by Orlicz function. *J. Math. Comput. Sci.* 4 (2)(2014), 267-277.
- [4] Ebadullah, K. DIFFERENCE SEQUENCE SPACES OF INVARIANT MEAN DEFINED BY ORLICZ FUNCTION. *Int. J. Math. Sci. Eng. Appl.* 6 (6)(2012), 101-110.
- [5] Ebadullah, K. On Certain Class of Difference Sequence Spaces. *Glob. J. Sci. Frontier Res. Math. Decision Sci.* 12 (11)(2012 ), 59-68.
- [6] Ebadullah, K.  $\sigma$ - Convergent Sequence Spaces Defined by Orlicz Function. *J. Math. Comput. Sci.* 6(6) (2016), 965-974.
- [7] Ebadullah, K.  $\sigma$ - Convergent Difference Sequence Spaces Defined by Orlicz Function. *J. Math. Comput. Sci.* 7 (1)(2017) 1-11
- [8] Ebadullah, K.  $\sigma$ - Convergent Difference Sequence Spaces Of Second Order Defined by Orlicz Function. *J. Math. Comput. Sci.* 7 (2) (2017) 237-248.
- [9] Khan, V.A., Ebadullah, K. *On a new difference sequence space of invariant mean defined by Orlicz functions.* *Bull. Allahabad Math. Soc.* 26(2)(2011), 259-272.

- [10] Kizmaz, H. On Certain sequence spaces, *Canad. Math. Bull.* 24(1981), 169-176.
- [11] Lindenstrauss, J. Tzafriri, L. On Orlicz sequence spaces. *Israel J. Math.*, 101(1971), 379-390.
- [12] Lorentz, C.G. A contribution to the theory of divergent series. *Acta Math.*, 80(1948), 167-190.
- [13] Maddox, I.J. *Elements of Functional Analysis*, Cambridge University Press. (1970)
- [14] Maddox, I.J. Some properties of paranormed sequence spaces., *J. London Math. Soc.* 1(1969), 316-322.
- [15] Mikael. *On some Difference Sequence Spaces*. Dogra - Tr. J. Math. 17 (1993), 18-24.
- [16] Mikael and Colak, R. On some generalized difference sequence spaces. *Soochow J. Math.* 21(4)(1995), 377-386.
- [17] Mursaleen, M. Matrix transformation between some new sequence spaces. *Houston J. Math.*, 9(1983), 505-509.
- [18] Mursaleen, M. On some new invariant matrix methods of summability. *Quart. J. Math. Oxford*, (2)34(1983), 77-86.
- [19] Musielak, J. Orlicz spaces and Modular spaces. *Lecture notes in Math.*(Springer-Verlag)1034:(1983).
- [20] Raimi, R.A. Invariant means and invariant matrix methods of summability. *Duke J. Math.*, 30(1963), 81-94.
- [21] Schafer, P. Infinite matrices and Invariant means. *Proc. Amer. Math. Soc.* 36 (1972), 104-110.