



Available online at <http://scik.org>

J. Math. Comput. Sci. 9 (2019), No. 6, 726-738

<https://doi.org/10.28919/jmcs/4227>

ISSN: 1927-5307

ON THE APPLICATIONS OF ALEPH (\aleph)-FUNCTION IN A SLIGHTLY DIFFERENT TYPE-2 BETA DENSITY MODEL

YASHWANT SINGH

Government College, Kaladera, Jaipur, Rajasthan, India

Copyright © 2019 the author(s). This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. In the present paper, the author has studied about the structures which are the products and ratios of statistically independently distributed positive real scalar random variables. The author has derived the exact density of a slightly different Type-2 beta density by the Mellin Transform and Hankel Transform of the unknown density and after that the unknown density has been derived in terms of Aleph- functions by taking the inverse Mellin transform and Inverse Hankel Transform. A more general structure of Type-2 beta density has also been discussed. Some special cases in terms of I -function are also given.

Key words: type-2 Beta density; aleph-Function; I -function; H -function; Mellin transform; inverse Mellin transform; Hankel transform; inverse Hankel transform.

2010 Mathematics Subject Classification: 33C10, 44A15.

1. INTRODUCTION

The I -function introduced by Saxena [4] will be represented and defined in slightly different manner as follows:

E-mail address: dryashu23@yahoo.in

Received July 16, 2019

$$I(z) = I_{p_i, q_i; R}^{m, n}(z) = I_{p_i, q_i; R}^{m, n} \left[z \left| \begin{matrix} (a_j, \alpha_j)_{1, n}, (a_{ji}, \alpha_{ji})_{n+1, p_i} \\ (b_j, \beta_j)_{1, m}, (b_{ji}, \beta_{ji})_{m+1, q_i} \end{matrix} \right. \right] = \frac{1}{2\pi i} \int_L \theta(s) z^{-s} ds \quad (1.1)$$

Where $i = \sqrt{-1}$, $z \neq 0$ and $z^{-s} = \exp[-\sin |z| + i \arg z]$ where $|z|$ represents the natural logarithm of $|z|$ and $\arg z$ is not the principal value. Here

$$\theta(s) = \frac{\prod_{j=1}^m \Gamma(b_j + B_j s) \prod_{j=1}^n \Gamma(1 - a_j - A_j s)}{\sum_{i=1}^R \left\{ \prod_{j=m+1}^q \Gamma(1 - b_{ji} - B_{ji} s) \prod_{j=n+1}^p \Gamma(a_{ji} + A_{ji} s) \right\}} \quad (1.2)$$

For $R = 1$, the I -function reduces to the H -function.

The \aleph -function introduced by Suland et.al. [6] defined and represented in the following form:

$$\begin{aligned} \aleph[z] &= \aleph_{p_i, q_i; \tau_i; r}^{m, n}[z] = \aleph_{p_i, q_i; \tau_i; r}^{m, n} \left[z \left| \begin{matrix} (a_j, \alpha_j)_{1, n}, [\tau_i(a_{ji}, \alpha_{ji})]_{n+1, p_i} \\ (b_j, \beta_j)_{1, m}, [\tau_i(b_{ji}, \beta_{ji})]_{m+1, q_i} \end{matrix} \right. \right] \\ &= \frac{1}{2\pi\omega} \int_L \theta(s) z^s ds \end{aligned} \quad (1.3)$$

where $\omega = \sqrt{-1}$;

$$\theta(s) = \frac{\prod_{j=1}^m \Gamma(b_j - \beta_j s) \prod_{j=1}^n \Gamma(1 - a_j + \alpha_j s)}{\sum_{i=1}^r \tau_i \left\{ \prod_{j=m+1}^{q_i} \Gamma(1 - b_{ji} + \beta_{ji} s) \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \alpha_{ji} s) \right\}} \quad (1.4)$$

We shall use the following notation:

$$A^* = (a_j, \alpha_j)_{1, n}, [\tau_i(a_{ji}, \alpha_{ji})]_{n+1, p_i}, B^* = (b_j, \beta_j)_{1, m}, [\tau_i(b_{ji}, \beta_{ji})]_{m+1, q_i}$$

The Mellin transform of $f(x)$ denoted by $M\{f(x); s\}$ or $F(s)$ is given by

$$M\{f(x); s\} = \int_0^{\infty} x^{s-1} f(x) dx \quad (1.5)$$

The Hankel transform of $f(x)$ denoted by $H_\nu\{f(x); p\}$ or $F_\nu(p)$ is given by

$$H_\nu\{f(x); p\} = \int_0^{\infty} x J_\nu(px) f(x) dx \quad (1.6)$$

A real scalar random variable x is said to have a real type-1 generalized beta distribution, if the density is of the following form ([1], p.121, eq. (4.8)):

$$f(x) = \begin{cases} \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1}; 0 < x < 1, \alpha > 0, \beta > 0 \\ 0, \text{elsewhere} \end{cases} \quad (1.7)$$

where the parameters α and β are real. The following discussion holds even when α and β are complex quantities. In this case, the conditions become $\text{Re}(\alpha) > 0, \text{Re}(\beta) > 0$ where $\text{Re}(\cdot)$ means the real part of (\cdot) .

A real scalar random variable x is said to have a real type-2 generalized beta distribution, if the density is of the following form ([1], p.121, eq. (4.8)):

$$f(x) = \begin{cases} \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1+x)^{-(\alpha+\beta)}; 0 < x < \infty, \alpha > 0, \beta > 0 \\ 0, \text{elsewhere} \end{cases} \quad (1.8)$$

Where the parameters α and β are real. The following discussion holds even when α and β are complex quantities. In this case, the conditions become $\text{Re}(\alpha) > 0, \text{Re}(\beta) > 0$ where $\text{Re}(\cdot)$ means the real part of (\cdot) .

2. GENERAL STRUCTURES

A real scalar random variable x is said to have a real generalized type-2 generalized beta distribution, if the density is of the following form:

$$f(x) = \begin{cases} \frac{\mathcal{A} \left(\frac{\alpha+tm}{\gamma} + \beta \right) A^{\frac{\alpha+tm}{\gamma}}}{\Gamma \left(\frac{\alpha+tm}{\gamma} \right) \Gamma(\beta)} x^{\alpha-1} (1+Ax)^{-(\alpha+\beta)} {}_2F_1(a, b; c; w; \mu; px') \\ 0, \text{elsewhere} \end{cases} \quad (2.1)$$

For $0 < x < \infty, \alpha > 0, \beta > 0, A > 0, a > 0, b > 0, c > 0, 1 + Ax^\gamma > 0$.

Where the parameters α and β are real. The following discussion holds even when α and β are complex quantities. In this case, the conditions become $\text{Re}(\alpha) > 0, \text{Re}(\beta) > 0$ where $\text{Re}(\cdot)$ means the real part of (\cdot) .

The h^{th} -moment of x , when x has the density in (2.1), is given by

$$E(x^h) = \frac{\Gamma \left(\frac{\alpha + tm + h}{\beta} \right)}{A^{\frac{h}{\beta}} \Gamma \left(\frac{\alpha + tm}{\beta} \right)} \quad (2.2)$$

For $\text{Re}(\alpha + tm + h) > 0, \beta > 0$.

When α and h are real, the moments can exist for some values of h also such that $\alpha + h > 0$.

The Mellin transform of $f(x)$ is obtained from (2.2) as:

$$M\{f(x)\} = \frac{\Gamma\left(\frac{\alpha + tm + s - 1}{\beta}\right)}{A^{\frac{s-1}{\beta}} \Gamma\left(\frac{\alpha + tm}{\beta}\right)} \quad (2.3)$$

For $\text{Re}(\alpha + tm + s - 1) > 0, s = v + 2r + 2 > 0, \beta > 0$.

The unknown density $f(x)$ is obtained in terms of \aleph -function by taking the inverse Mellin transform of (2.3). That is

$$f(x) = \frac{A^{\frac{1}{\beta}}}{\Gamma\left(\frac{\alpha + tm}{\beta}\right)} \aleph_{0,1;\tau_j;R}^{s,1,0} \left[x \left| \begin{matrix} - \\ \left(\frac{\alpha + tm - 1}{\beta}, \frac{1}{\beta}\right) \end{matrix} \right. \right] \quad (2.4)$$

The Hankel transform of $f(x)$ is obtained from (2.2) as:

$$H\{f(x)\} = J_\nu(p) \frac{\Gamma\left(\frac{\alpha + tm + s - 1}{\beta}\right)}{A^{\frac{s-1}{\beta}} \Gamma\left(\frac{\alpha + tm}{\beta}\right)} \quad (2.5)$$

For $\text{Re}(\alpha + tm + s - 1) > 0, s = v + 2r + 2 > 0, \beta > 0$.

The unknown density $f(x)$ is obtained in terms of \aleph -function by taking the inverse Hankel transform of (2.5). That is

$$f(x) = J_\nu(p) \frac{A^{\frac{1}{\beta}}}{\Gamma\left(\frac{\alpha + tm}{\gamma}\right)} \aleph_{0,1;\tau_j;R}^{s,1,0} \left[A^{\frac{1}{\beta}} x \left| \begin{matrix} - \\ \left(\frac{\alpha + tm - 1}{\beta}, \frac{1}{\beta}\right) \end{matrix} \right. \right] \quad (2.6)$$

Consider a set of real scalar random variables x_1, \dots, x_k , mutually independently distributed, where x_j has the density in (2.1) with the parameters $\alpha_j, \beta_j, A_j; j = 1, \dots, k$ and consider the product

$$u = x_1 x_2 \dots x_k \quad (2.7)$$

In the standard terminology in statistical literature, the h^{th} moment of u , when u has the density in (2.1), is given by

$$E(u) = \frac{\Gamma\left(\frac{\alpha_j + tm + h}{\beta_j}\right)}{A^{\frac{h}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \quad (2.8)$$

For $\text{Re}(\alpha_j + tm + h) > 0, \beta_j > 0; j = 1, \dots, k$

Then the Mellin transform of $g(u)$ of u is obtained from the property of the statistical independent and is given by

$$M[g(u)] = M[x_1^{s-1}] \dots M[x_k^{s-1}] \quad (2.9)$$

$$M[g(u)] = \prod_{j=1}^k \frac{\Gamma\left(\frac{\alpha_j + tm + s - 1}{\beta_j}\right)}{A^{\frac{s-1}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \quad (2.10)$$

For $\text{Re}(\alpha_j + tm + s - 1) > 0, \beta_j > 0, s = v + 2r + 2 > 0$

The unknown density $f(x)$ is obtained in terms of \aleph -function by taking the inverse Mellin transform of (2.10). That is

$$g(u) = \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \aleph_{0,k;\tau_j;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{1}{\beta_j}} u \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, k \right] \quad (2.11)$$

Then the Hankel transform of $g(u)$ of u is obtained from the property of the statistical independent and is given by:

$$H[g(u)] = H[x_1 J_\nu(px_1)] H[x_2 J_\nu(px_2)] \dots H[x_k J_\nu(px_k)] \quad (2.12)$$

$$H\{g(u)\} = J_\nu(p) \prod_{j=1}^k \frac{\Gamma\left(\frac{\alpha_j + tm + s - 1}{\beta_j}\right)}{A^{\frac{s-1}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \quad (2.13)$$

For $\text{Re}(\alpha_j + tm + s - 1) > 0, \beta_j > 0, s = v + 2r + 2 > 0$

The unknown density $f(x)$ is obtained in terms of \aleph -function by taking the inverse Hankel transform of (2.13). That is

$$g(u) = J_\nu(p) \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \aleph_{0,k;\tau_j;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{1}{\beta_j}} u \left| \begin{array}{c} \frac{1}{\beta_j} \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{array} \right. ; j = 1, \dots, k \right] \quad (2.14)$$

If we consider more general structures in the same category. For example, consider the structure $u_1 = x_1^{\delta_1} x_2^{\delta_2} \dots x_k^{\delta_k}$, $\delta_j > 0$, $j = 1, \dots, k$ (2.15)

Where x_1, \dots, x_k are mutually independently distributed as in (2.1).

Then the Mellin transform of $g(u_1)$ of u_1 is given as

$$M[g(u_1)] = M[x_1^{\gamma_1(s-1)}] \dots M[x_k^{\gamma_k(s-1)}] \quad (2.16)$$

$$M[g(u_1)] = \prod_{j=1}^k \frac{\Gamma\left(\frac{\alpha_j + tm + \delta_j(s-1)}{\beta_j}\right)}{A^{\frac{\delta_j(s-1)}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \quad (2.17)$$

For $\text{Re}(\alpha_j + tm + s - 1) > 0$, $s = \nu + 2r + 2$, $\delta_j > 0$.

The unknown density $g(u_1)$ is obtained in terms of \aleph -function by taking the inverse Mellin transform of (2.17). That is

$$g(u_1) = \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \aleph_{0,k;\tau_j;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{\delta_j}{\beta_j}} u_1 \left| \begin{array}{c} \frac{\delta_j}{\beta_j} \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{array} \right. ; j = 1, \dots, k \right] \quad (2.18)$$

For $\text{Re}(\alpha_j + tm + s - 1) > 0$, $s = \nu + 2r + 2$, $\delta_j > 0$.

The Hankel transform of $g(u_1)$ of u_1 is obtained from the property of the statistical independent and is given by:

$$H[g(u_1)] = H[x_1^{\delta_1} J_\nu(px_1^{\delta_1})] H[x_2^{\delta_2} J_\nu(px_2^{\delta_2})] \dots H[x_k^{\delta_k} J_\nu(px_k^{\delta_k})] \quad (2.19)$$

$$H\{g(u_1)\} = J_\nu(p) \prod_{j=1}^k \frac{\Gamma\left(\frac{\alpha_j + tm + \delta_j(s-1)}{\gamma_j}\right)}{A^{\frac{\delta_j(s-1)}{\gamma_j}} \Gamma\left(\frac{\alpha_j + tm}{\gamma_j}\right)} \quad (2.20)$$

For $\operatorname{Re}(\alpha_j + tm + s - 1) > 0, s = \nu + 2r + 2 > 0, \delta_j > 0$.

The unknown density $g(u_1)$ is obtained in terms of \aleph -function by taking the inverse Hankel transform of (2.20). That is

$$g(u_1) = J_\nu(p) \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \aleph_{0,k;\tau_i;R}^{s,k,0} \left[\prod_{j=1}^k A^{\frac{\delta_j}{\beta_j}} u_1 \left| \begin{array}{c} \text{---} \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{array} \right. ; j = 1, \dots, k \right] \quad (2.21)$$

For $\operatorname{Re}(\alpha_j + tm + s - 1) > 0, s = \nu + 2r + 2 > 0, \delta_j > 0$.

A More General Structure

We can consider more general structures. Let

$$w = \frac{x_1, x_2, \dots, x_r}{x_{r+1}, \dots, x_k} \quad (2.22)$$

Where x_1, \dots, x_k , mutually independently distributed real random variables having the density in (2.1) with x_j having parameters $\alpha_j, \beta_j, A_j; j = 1, \dots, k$.

Then the Mellin transform of $g(w)$ is given by

$$M[g(w)] = M[x_1^{s-1}] \dots M[x_r^{s-1}] M[x_{r+1}^{-(s-1)}] \dots M[x_k^{-(s-1)}] \quad (2.23)$$

$$M[g(w)] = \prod_{j=1}^k \frac{1}{A^{\frac{-1}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ \prod_{j=1}^r \frac{\Gamma\left(\frac{\alpha_j + tm + s - 1}{\beta_j}\right)}{A^{\frac{s}{\beta_j}}} \right\} \left\{ \prod_{j=r+1}^k \frac{\Gamma\left(\frac{\alpha_j + tm - (s-1)}{\beta_j}\right)}{A^{\frac{-s}{\beta_j}}} \right\} \quad (2.24)$$

For $\operatorname{Re}(\alpha_j + tm \pm (s-1)) > 0, s = \nu + 2r + 2 > 0$

The unknown density $g(w)$ is obtained in terms of \aleph -function by taking the inverse Mellin transform of (2.24). That is

$$g(w) = \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ \aleph_{0,r;\tau_j;R}^{s,r,0} \left[\prod_{j=1}^r A^{\frac{1}{\beta_j}} w \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, r \right] \right\} \\ \left\{ \aleph_{0,k-r;\tau_j;R}^{s,k-r,0} \left[\prod_{j=r+1}^k A^{\frac{1}{\beta_j}} w \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + 1}{\beta_j}, \frac{1}{\beta_j}\right) \\ - \end{matrix} \right. ; j = r + 1, \dots, k \right] \right\} \quad (2.25)$$

Then the Hankel transform of $g(w)$ is given as:

$$H[g(w)] = H[x_1 J_\nu(px_1)] \dots H[x_r J_\nu(px_r)] \\ H[x_{r+1}^{-1} J_\nu(px_{r+1}^{-1})] \dots H[x_k^{-1} J_\nu(px_k^{-1})] \quad (2.26)$$

$$H\{g(w)\} = J_\nu(p) \prod_{j=1}^k \frac{1}{A^{\frac{-1}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ \prod_{j=1}^r \frac{\Gamma\left(\frac{\alpha_j + tm + s - 1}{\beta_j}\right)}{A^{\frac{s}{\beta_j}}} \right\} \\ \left\{ \prod_{j=r+1}^k \frac{\Gamma\left(\frac{\alpha_j + tm - (s - 1)}{\beta_j}\right)}{A^{\frac{-s}{\beta_j}}} \right\} \quad (2.27)$$

For $\text{Re}(\alpha_j + tm \pm (s - 1)) > 0, s = \nu + 2r + 2 > 0$

The unknown density $g(w)$ is obtained in terms of \aleph -function by taking the inverse Hankel transform of (2.27). That is

$$g(w) = J_\nu(p) \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ \aleph_{0,r;\tau_j;R}^{s,r,0} \left[\prod_{j=1}^r A^{\frac{1}{\beta_j}} w \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, r \right] \right\} \\ \left\{ \aleph_{k-r,0;\tau_j;R}^{s,0,k-r} \left[\prod_{j=r+1}^k A^{\frac{1}{\beta_j}} w \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + 1}{\beta_j}, \frac{1}{\beta_j}\right) \\ - \end{matrix} \right. ; j = r + 1, \dots, k \right] \right\} \quad (2.28)$$

We can consider more general structures in the same category. For example, consider the structure

$$w_1 = \frac{x_1^{\delta_1}, \dots, x_r^{\delta_r}}{x_{r+1}^{\delta_{r+1}}, \dots, x_k^{\delta_k}} \quad (2.29)$$

where x_1, \dots, x_k , mutually independently distributed real random variables having the density in (2.1) with x_j having parameters $\alpha_j, \beta_j; j=1, \dots, k$.

Then the Mellin transform of $g(w_1)$ is given by

$$M[g(w_1)] = M[x_1^{\delta_1(s-1)}] \dots M[x_r^{\delta_r(s-1)}] M[x_{r+1}^{-\delta_{r+1}(s-1)}] \dots M[x_k^{-\delta_k(s-1)}] \quad (2.30)$$

$$M[g(w_1)] = \prod_{j=1}^k \frac{1}{A^{\frac{-\delta_j}{\beta_j}} \Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ \prod_{j=1}^r \frac{\Gamma\left(\frac{\alpha_j + tm + \delta_j(s-1)}{\beta_j}\right)}{A^{\frac{\delta_j s}{\beta_j}}} \right\} \left\{ \prod_{j=r+1}^k \frac{\Gamma\left(\frac{\alpha_j + tm - \delta_j(s-1)}{\beta_j}\right)}{A^{\frac{-s\delta_j}{\beta_j}}} \right\} \quad (2.31)$$

For $\text{Re}(\alpha_j + tm \pm (s-1)) > 0, s = v + 2r + 2 > 0, \delta_j > 0$.

The unknown density $g(w_1)$ is obtained in terms of \aleph -function by taking the inverse Mellin transform of (2.31). That is

$$g(w_1) = \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ \aleph_{0,r;\tau_j;R}^{r,0} \left[\prod_{j=1}^r A^{\frac{\delta_j}{\beta_j}} w_1 \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{matrix} \right. ; j=1, \dots, r \right] \right\} \left\{ \aleph_{k-r,0;\tau_j;R}^{0,k-r} \left[\prod_{j=r+1}^k A^{\frac{-\delta_j}{\beta_j}} w_1 \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \\ - \end{matrix} \right. ; j=r+1, \dots, k \right] \right\} \quad (2.32)$$

For $\text{Re}(\alpha_j + tm \pm (s-1)) > 0, s = v + 2r + 2 > 0, \delta_j > 0$.

Then the Hankel transform of $g(w_1)$ is given as:

$$H[g(w_1)] = H[x_1^{\delta_1} J_\nu(px_1^{\delta_1})] \dots H[x_r^{\delta_r} J_\nu(px_r^{\delta_r})]$$

$$H \left[x_{r+1}^{-\delta_{r+1}} J_\nu \left(p x_{r+1}^{-\delta_{r+1}} \right) \right] \dots H \left[x_k^{-\delta_k} J_\nu \left(p x_k^{-\delta_k} \right) \right] \quad (2.33)$$

$$H \{ g(w_1) \} = J_\nu(p) \prod_{j=1}^k \frac{1}{A^{\frac{-\delta_j}{\beta_j}} \Gamma \left(\frac{\alpha_j + tm}{\beta_j} \right)} \left\{ \prod_{j=1}^r \frac{\Gamma \left(\frac{\alpha_j + tm + \delta_j (s-1)}{\beta_j} \right)}{A^{\frac{\delta_j s}{\beta_j}}} \right\} \left\{ \prod_{j=r+1}^k \frac{\Gamma \left(\frac{\alpha_j + tm - \delta_j (s-1)}{\beta_j} \right)}{A^{\frac{-s\delta_j}{\beta_j}}} \right\} \quad (2.34)$$

For $\operatorname{Re}(\alpha_j + tm \pm (s-1)) > 0, \delta_j > 0, s = \nu + 2r + 2 > 0$

The unknown density $g(w_1)$ is obtained in terms of \aleph -function by taking the inverse Hankel transform of (2.34). That is

$$g(w_1) = J_\nu(p) \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma \left(\frac{\alpha_j + tm}{\beta_j} \right)} \left\{ \aleph_{0,r;\tau_i;R}^{s,r,0} \left[\prod_{j=1}^r A^{\frac{\delta_j}{\beta_j}} w_1 \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j} \right) \end{matrix} \right. ; j = 1, \dots, r \right] \right\} \left\{ \aleph_{k-r,0;\tau_i;R}^{s,0,k-r} \left[\prod_{j=r+1}^k A^{\frac{-\delta_j}{\beta_j}} w_1 \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j} \right) \\ - \end{matrix} \right. ; j = r+1, \dots, k \right] \right\} \quad (2.35)$$

For $\operatorname{Re}(\alpha_j + tm \pm (s-1)) > 0, \delta_j > 0, s = \nu + 2r + 2 > 0$.

3. SPECIAL CASES

If we take $\tau_i = 1$ in (2.4), the unknown density $f(x)$ is obtained in terms of I -function. That is

$$f(x) = \frac{A^{\frac{1}{\beta}}}{\Gamma \left(\frac{\alpha + tm}{\beta} \right)} I_{0,1;R}^{1,0} \left[A^{\frac{1}{\beta}} x \left| \begin{matrix} - \\ \left(\frac{\alpha + tm - 1}{\beta}, \frac{1}{\beta} \right) \end{matrix} \right. \right] \quad (3.1)$$

For $R=1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.6), the unknown density $f(x)$ is obtained in terms of I -function. That is

$$f(x) = J_v(p) \frac{A^{\frac{1}{\beta}}}{\Gamma\left(\frac{\alpha + tm}{\beta}\right)} I_{0,1;R}^{1,0} \left[A^{\frac{1}{\beta}} x \left| \begin{matrix} - \\ \left(\frac{\alpha + tm - 1}{\beta}, \frac{1}{\beta}\right) \end{matrix} \right. \right] \quad (3.2)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.11), the unknown density $g(u)$ is obtained in terms of I -function. That is

$$g(u) = \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} I_{0,k;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{1}{\beta_j}} u \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, k \right] \quad (3.3)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.14), the unknown density $g(u)$ is obtained in terms of I -function. That is

$$g(u) = J_v(p) \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} I_{0,k;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{1}{\beta_j}} u \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, k \right] \quad (3.4)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.18), the unknown density $g(u_1)$ is obtained in terms of I -function. That is

$$g(u_1) = \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} I_{0,k;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{\delta_j}{\beta_j}} u_1 \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, k \right] \quad (3.5)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.21), the unknown density $g(u_1)$ is obtained in terms of I -function. That is

$$g(u_1) = J_v(p) \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} I_{0,k;R}^{k,0} \left[\prod_{j=1}^k A^{\frac{\delta_j}{\beta_j}} u_1 \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{matrix} \right. \right] \quad (3.6)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.25), the unknown density $g(w)$ is obtained in terms of I -function. That is

$$g(w) = \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ I_{0,r;R}^{r,0} \left[\prod_{j=1}^r A^{\frac{1}{\beta_j}} w \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, r \right] \right. \\ \left. \left\{ I_{k-r,0;R}^{0,k-r} \left[\prod_{j=r+1}^k A^{\frac{1}{\beta_j}} w \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + 1}{\beta_j}, \frac{1}{\beta_j}\right) \\ - \end{matrix} \right. ; j = r + 1, \dots, k \right] \right\} \right. \quad (3.7)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.28), the unknown density $g(w)$ is obtained in terms of I -function. That is

$$g(w) = J_v(p) \prod_{j=1}^k \frac{A^{\frac{1}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ I_{0,r;R}^{r,0} \left[\prod_{j=1}^r A^{\frac{1}{\beta_j}} w \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - 1}{\beta_j}, \frac{1}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, r \right] \right\} \\ \left\{ I_{k-r,0;R}^{0,k-r} \left[\prod_{j=r+1}^k A^{\frac{1}{\beta_j}} w \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + 1}{\beta_j}, \frac{1}{\beta_j}\right) \\ - \end{matrix} \right. ; j = r + 1, \dots, k \right] \right\} \quad (3.8)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.32), the unknown density $g(w_1)$ is obtained in terms of I -function. That is

$$g(w_1) = \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ I_{0,r;R}^{r,0} \left[\prod_{j=1}^r A^{\frac{\delta_j}{\beta_j}} w_1 \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, r \right] \right\} \\ \left\{ I_{k-r,0;R}^{0,k-r} \left[\prod_{j=r+1}^k A^{\frac{-\delta_j}{\beta_j}} w_1 \left| \begin{matrix} \left(1 - \frac{\alpha_j + tm + \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \\ - \end{matrix} \right. ; j = r + 1, \dots, k \right] \right\} \quad (3.9)$$

For $R = 1$, the I -function reduces to the H -function.

If we take $\tau_i = 1$ in (2.35), the unknown density $g(w_1)$ is obtained in terms of I -function. That is

$$g(w_1) = J_v(p) \prod_{j=1}^k \frac{A^{\frac{\delta_j}{\beta_j}}}{\Gamma\left(\frac{\alpha_j + tm}{\beta_j}\right)} \left\{ I_{0,r;R}^{r,0} \left[\prod_{j=1}^r A^{\frac{\delta_j}{\beta_j}} w_1 \left| \begin{matrix} - \\ \left(\frac{\alpha_j + tm - \delta_j}{\beta_j}, \frac{\delta_j}{\beta_j}\right) \end{matrix} \right. ; j = 1, \dots, r \right] \right\}$$

$$\left\{ I_{k-r,0;R}^{0,k-r} \left[\prod_{j=r+1}^k A_{\beta_j}^{-\delta_j} w_1 \left(\begin{matrix} - \\ \left(1 - \frac{\alpha_j + \delta_j}{\beta_j}, \delta_j \right) \end{matrix} \right); j = r+1, \dots, k \right] \right\} \quad (3.10)$$

For $R = 1$, the I -function reduces to the H -function.

Conflict of Interests

The author(s) declare that there is no conflict of interests.

REFERENCES

- [1] Mathai, A.M.; A pathway to matrix-variate gamma and normal densities, *Linear Alg. Appl.* 396 (2005), 317-328.
- [2] Mathai, A.M. and Saxena, R.K.; *The H-function with Applications in Statistics and Other Disciplines*, Wiley Eastern, New Delhi and Wiley Halsted, New York (1978).
- [3] Mathai, A.M., Saxena, R.K. and Haubold, H.J.; *The H-function: Theory and Applications*, CRC Press, New York (2010).
- [4] Saxena, V.P.; Formal solution of certain new pair of dual integral equations involving H -functions, *Proc. Nat. Acad. India, Sect. A* 52(1982), 366-375.
- [5] Srivastava, H.M., Gupta, K.C. and Goyal, S.P.; *The H-function of One and Two Variables with Applications*, South Asian Publisher, New Delhi, (1982).
- [6] Sudland, N., Baumann, B. and Nonnenmacher, T.F.; Open problem: who knows about the Aleph (\aleph)-functions? *Frac. Calc. Appl. Annl.* 1(4)(1998), 401-402.