



Available online at <http://scik.org>

J. Math. Comput. Sci. 10 (2020), No. 6, 2871-2890

<https://doi.org/10.28919/jmcs/4910>

ISSN: 1927-5307

A NEW IMPULSIVE SEQUENTIAL MULTI-ORDERS FRACTIONAL DIFFERENTIAL EQUATION WITH BOUNDARY CONDITIONS

AMER DABABNEH, B. SAMI, MA'MON ABU HAMMAD*, AMJED ZRAIQAT

Department of Mathematics, Al-Zaytoonah University of Jordan, Jordan

Copyright © 2020 the author(s). This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. In this paper, we present a new impulsive sequential multi-orders fractional differential equation is studied. The existence and uniqueness results are obtained for a nonlinear problem with fractional integral boundary conditions applying standard fixed point theorems. An example for the illustration of the main result is given.

Keywords: impulsive sequential fractional differential equations; boundary value problem; Banach fixed point theorem.

2010 AMS Subject Classification: 34A08.

1. INTRODUCTION

Nowadays, fractional differential equations have attracted a lot of attention due to its wide range of applications in many practical problems such as in physics, engineering, economics, and so on; see [1–5].

Impulsive sequential differential equations have extensively been studied in the past two decades. Indeed impulsive differential equations are used to describe the dynamics of processes in which sudden, discontinuous jumps occur. Such processes are naturally seen in harvesting, earthquakes, diseases, and so forth. Recently, fractional impulsive differential equations have

*Corresponding author

E-mail address: mabuhammad@gmail.com

Received August 4, 2020

attracted the attention of many researchers. For the general theory and applications of such equations we refer the interested reader to see [6] and [7–20] and the references therein. Based on previous studies, in this topic we concentrate on the existence results of solutions of the following problem:

$$(1) \quad ({}^cD_{t_k^+}^{\beta_k} + \lambda {}^cD_{t_k^+}^{\beta_k-1})x(t) = f(t, x(t)), \quad 1 < \beta_k \leq 2, k = 0, 1, \dots, q; t \in J',$$

$$\Delta x(t_k) = \psi_k(x(t_k)), \quad \Delta x'(t_k) = \psi_k^*(x(t_k)), \quad k = 1, \dots, q,$$

$$x(0) = \sum_{k=0}^q \lambda_k I_{t_k^+}^{\alpha_k} x(\eta_k), \quad x'(0) = 0.$$

where ${}^cD_{t_k^+}^{\beta_k}$ is the Caputo fractional derivative of order $\beta_k \in (1, 2]$ and $I_{t_k^+}^{\alpha_k}$ is fractional Riemann-Liouville integral of order $\alpha_k > 0$, $f \in (J \times \mathbb{R}, \mathbb{R})$, $\psi_k, \psi_k^* \in C(\mathbb{R} \times \mathbb{R})$, $\lambda_k \in \mathbb{R}$, $\lambda \in \mathbb{R}^+$, $J = [0, T]$, $J' = J \setminus \{t_1, \dots, t_q\}$, $0 = t_0 < t_1 < \dots < t_k \dots < t_q < t_{q+1} = T$, $\Delta x(t_k) = x(t_k^+) - x(t_k^-)$ and $\Delta x'(t_k) = x'(t_k^+) - x'(t_k^-)$. Here, respectively, the right and the left limits of $x(t)$ at $t = t_k^+$ are represented by $x(t_k^+)$ and $x(t_k^-)$.

2. BASIC MATERIALS

We introduce throughout this section preliminary facts that will be used in this paper. We fix $J_0 = [0, t_1]$, $J_{k-1} = (t_{k-1}, t_k]$, and $k = 1, 2, \dots, q+1$ with $t_{q+1} = T$ and define the Banach space

$$PC(J) = \{x : J \rightarrow \mathbb{R} \mid x \in C(J'), \text{ and } x(t_k^+), x(t_k^-) \text{ exist, and } x(t_k^+) = x(t_k), 1 \leq k \leq q\},$$

with the norm $\|x\|_{PC} = \sup \{|x(t)| : t \in J\}$.

Definition 2.1. *The fractional integral of order β with the lower limit zero for a function $f : [x, \infty) \rightarrow \mathbb{R}$ is defined as*

$$I_{x^+}^\beta f(t) = \frac{1}{\Gamma(\beta)} \int_x^t (t-s)^{\beta-1} f(s) ds, \quad t > 0, \beta > 0,$$

provided the right side is point-wise defined on $[x, \infty)$, where $\Gamma(\cdot)$ is the gamma function.

Definition 2.2. *The Caputo fractional derivative of order $\beta > 0$, of a function $f : [x, \infty) \rightarrow \mathbb{R}$ can be written as*

$$D_{x^+}^\beta f(t) = \frac{1}{\Gamma(n-\beta)} \left(\frac{d}{dt} \right)^n \int_x^t (t-s)^{n-\beta-1} f(s) ds, \quad t > 0, n-1 < \beta < n.$$

where $n = [\beta] + 1$ (the notation $[\beta]$ stands for the largest integer not greater than β).

Lemma 2.1. For a given $h \in PC[0, T]$ a function $x \in PC(J, \mathbb{R})$ is a solution of the impulsive sequential fractional differential equation

$$(2) \quad (^cD_{t_k^+}^{\beta_k} + \lambda ^cD_{t_k^+}^{\beta_k-1})u(t) = h(t), \quad 1 < \beta_k \leq 2, k = 0, 1, 2, \dots, q, t \in J',$$

$$(3) \quad \Delta x(t_k) = \psi_k(x(t_k)), \quad \Delta x'(t_k) = \psi_k^*(x(t_k)), \quad k = 1, \dots, q,$$

with boundary condition

$$(4) \quad x(0) = \sum_{k=0}^q \lambda_k I_{t_k^+}^{\alpha_k} x(\eta_k), \quad x'(0) = 0,$$

if and only if

$$(5) \quad x(t) = \begin{cases} \int_0^t e^{-\lambda(t-s)} I_{0^+}^{\beta_0-1} h(s) ds + \wp, & t \in J_0; \\ \int_{t_k}^t e^{-\lambda(t-s)} I_{t_k^+}^{\beta_k-1} h(s) ds \\ + \sum_{j=1}^k e^{-\lambda(t-t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} I_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds \right. \\ - \frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \left. \right] \\ + \sum_{j=1}^k \left[\frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\ + \wp, & t \in J_k, k = 1, \dots, q, \end{cases}$$

where

$$\begin{aligned} \wp &= \left(1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right)^{-1} \\ &\times \left\{ \sum_{k=0}^q \lambda_k I_{t_k^+}^{\alpha_k} \left(\int_{t_k}^{\eta_k} e^{-\lambda(r-s)} I_{t_k^+}^{\beta_k-1} h(s) ds \right) (\eta_k) \right. \\ &+ \sum_{k=0}^q \sum_{j=1}^k \lambda_k I_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k - t_j)} \\ &\left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} I_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds - \frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \right. \\ &\left. + \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \right\}. \end{aligned}$$

Proof. Assume that x is a solution of (1). For any $t \in J_0$, we have

$$(6) \quad x(t) = \int_0^t e^{-\lambda(t-s)} I_{0^+}^{\beta_0-1} h(s) ds + e^{-\lambda t} a_1 + a_2, \quad t \in J_0,$$

where a_1 and $a_2 \in \mathbb{R}$. Differentiating the obtained linear equation (6) on J_0 , leads to

$$(7) \quad x'(t) = -\lambda \int_0^t e^{-\lambda(t-s)} I_{0^+}^{\beta_0-1} h(s) ds + I_{0^+}^{\beta_0-1} h(t) - \lambda e^{-\lambda t} a_1.$$

If $t \in J_1$, then

$$(8) \quad \begin{aligned} x(t) &= \int_{t_1}^t e^{-\lambda(t-s)} I_{t_1^+}^{\beta_1-1} h(s) ds + e^{-\lambda(t-t_1)} b_1 + b_2, \\ x'(t) &= -\lambda \int_{t_1}^t e^{-\lambda(t-s)} I_{t_1^+}^{\beta_1-1} h(s) ds + I_{t_1^+}^{\beta_1-1} h(t) - \lambda e^{-\lambda(t-t_1)} b_1, \end{aligned}$$

for some $b_1, b_2 \in \mathbb{R}$. Thus,

$$(9) \quad \begin{aligned} x(t_1^-) &= \int_0^{t_1} e^{-\lambda(t_1-s)} I_{0^+}^{\beta_0-1} h(s) ds + e^{-\lambda t_1} a_1 + a_2, \\ x'(t_1^-) &= -\lambda \int_0^{t_1} e^{-\lambda(t_1-s)} I_{0^+}^{\beta_0-1} h(s) ds + I_{0^+}^{\beta_0-1} h(t_1) - \lambda e^{-\lambda t_1} a_1, \\ x(t_1^+) &= b_1 + b_2, \\ x'(t_1^+) &= -\lambda b_1. \end{aligned}$$

Now, by the following impulsive conditions

$$(10) \quad \Delta x(t_1) = x(t_1^+) - x(t_1^-) = \psi_1(x(t_1)),$$

$$\Delta x'(t_1) = x'(t_1^+) - x'(t_1^-) = \psi_1^*(x(t_1)),$$

we can get that

$$(11) \quad \begin{aligned} b_1 &= \int_0^{t_1} e^{-\lambda(t_1-s)} I_{0^+}^{\beta_0-1} h(s) ds - \frac{1}{\lambda} I_{0^+}^{\beta_0-1} h(t_1) + e^{-\lambda t_1} a_1 - \frac{1}{\lambda} \psi_1^*(x(t_1)), \\ b_2 &= \frac{1}{\lambda} I_{0^+}^{\beta_0-1} h(t_1) + \psi_1(x(t_1)) + \frac{1}{\lambda} \psi_1^*(x(t_1)) + a_2. \end{aligned}$$

Consequently,

$$\begin{aligned} x(t) = & \int_{t_1}^t e^{-\lambda(t-s)} \mathbf{I}_{t_1^+}^{\beta_1-1} h(s) ds + e^{-\lambda(t-t_1)} \left[\int_0^{t_1} e^{-\lambda(t_1-s)} \mathbf{I}_{0^+}^{\beta_0-1} h(s) ds \right. \\ & - \frac{1}{\lambda} \mathbf{I}_{0^+}^{\beta_0-1} h(t_1) - \frac{1}{\lambda} \psi_1^*(x(t_1)) \Big] + \left[\frac{1}{\lambda} \mathbf{I}_{0^+}^{\beta_0-1} h(t_1) \right. \\ & \left. + \psi_1(x(t_1)) + \frac{1}{\lambda} \psi_1^*(x(t_1)) \right] + e^{-\lambda t} a_1 + a_2, t \in J_1. \end{aligned}$$

If $t \in J_2$, then

$$\begin{aligned} (12) \quad x(t) = & \int_{t_2}^t e^{-\lambda(t-s)} \mathbf{I}_{t_2^+}^{\beta_2-1} h(s) ds + e^{-\lambda(t-t_2)} c_1 + c_2, \\ x'(t) = & -\lambda \int_{t_2}^t e^{-\lambda(t-s)} \mathbf{I}_{t_2^+}^{\beta_2-1} h(s) ds + \mathbf{I}_{t_2^+}^{\beta_1-1} h(t) - \lambda e^{-\lambda(t-t_2)} c_1, \end{aligned}$$

for some $c_1, c_2 \in \mathbb{R}$. Thus,

$$\begin{aligned} (13) \quad x(t_2^-) = & \int_{t_1}^{t_2} e^{-\lambda(t_2-s)} \mathbf{I}_{t_1^+}^{\beta_1-1} h(s) ds + e^{-\lambda(t_2-t_1)} b_1 + b_2, \\ x'(t_2^-) = & -\lambda \int_{t_1}^{t_2} e^{-\lambda(t_2-s)} \mathbf{I}_{t_1^+}^{\beta_1-1} h(s) ds + \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) - \lambda e^{-\lambda(t_2-t_1)} b_1, \\ x(t_2^+) = & c_1 + c_2, \\ x'(t_2^+) = & -\lambda c_1. \end{aligned}$$

Now, by the following impulsive conditions

$$\begin{aligned} (14) \quad \Delta x(t_2) = & x(t_2^+) - x(t_2^-) = \psi_2(x(t_2)), \\ \Delta x'(t_2) = & x'(t_2^+) - x'(t_2^-) = \psi_2^*(x(t_2)), \end{aligned}$$

we can get that

$$\begin{aligned} (15) \quad c_1 = & \int_{t_1}^{t_2} e^{-\lambda(t_2-s)} \mathbf{I}_{t_1^+}^{\beta_1-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) + e^{-\lambda(t_2-t_1)} b_1 - \frac{1}{\lambda} \psi_2^*(x(t_2)), \\ c_2 = & \frac{1}{\lambda} \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) + \psi_2(x(t_2)) + \frac{1}{\lambda} \psi_2^*(x(t_2)) + b_2. \end{aligned}$$

Consequently,

$$\begin{aligned}
 (16) \quad x(t) &= \int_{t_2}^t e^{-\lambda(t-s)} \mathbf{I}_{t_2^+}^{\beta_2-1} h(s) ds + e^{-\lambda(t-t_2)} \\
 &\times \left[\int_{t_1}^{t_2} e^{-\lambda(t_2-s)} \mathbf{I}_{t_1^+}^{\beta_1-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) \right. \\
 &\left. - \frac{1}{\lambda} \psi_2^*(x(t_2)) \right] + \frac{1}{\lambda} \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) + \psi_2(x(t_2)) \\
 &+ \frac{1}{\lambda} \psi_2^*(x(t_2)) + e^{-\lambda(t-t_1)} b_1 + b_2 \\
 &= \int_{t_2}^t e^{-\lambda(t-s)} \mathbf{I}_{t_2^+}^{\beta_2-1} h(s) ds + e^{-\lambda(t-t_2)} \\
 &\left[\int_{t_1}^{t_2} e^{-\lambda(t_2-s)} \mathbf{I}_{t_1^+}^{\beta_1-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) \right. \\
 &\left. - \frac{1}{\lambda} \psi_2^*(x(t_2)) \right] + \frac{1}{\lambda} \mathbf{I}_{t_1^+}^{\beta_1-1} h(t_2) + \psi_2(x(t_2)) + \frac{1}{\lambda} \psi_2^*(x(t_2)) \\
 &+ e^{-\lambda(t-t_1)} \left[\int_0^{t_1} e^{-\lambda(t_1-s)} \mathbf{I}_{0^+}^{\beta_0-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{0^+}^{\beta_0-1} h(t_1) - \frac{1}{\lambda} \psi_1^*(x(t_1)) \right] \\
 &+ \frac{1}{\lambda} \mathbf{I}_{0^+}^{\beta_0-1} h(t_1) + \psi_1(x(t_1)) + \frac{1}{\lambda} \psi_1^*(x(t_1)) + e^{-\lambda t} a_1 + a_2, \quad t \in J_2.
 \end{aligned}$$

Repeating the process in this way, we get

$$\begin{aligned}
 (17) \quad x(t) &= \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds + \sum_{j=1}^k e^{-\lambda(t-t_j)} \\
 &\times \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\
 &+ \sum_{j=1}^k \left[\frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] + e^{-\lambda t} a_1 + a_2, \\
 t \in J_k, k &= 1, 2, \dots, q.
 \end{aligned}$$

Taking (6), (7) and (17) to the boundary conditions,

$$x(0) = \sum_{k=0}^q \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} x(\eta_k), \quad x'(0) = 0,$$

implies $a_1 = 0$. For $t \in J_k$, we have

$$\begin{aligned} \mathbf{I}_{t_k^+}^{\alpha_k} x(t) &= \mathbf{I}_{t_k^+}^{\alpha_k} \left(\int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_{k-1}-1} h(s) ds \right) (t) + \sum_{j=1}^k \mathbf{I}_{t_k^+}^{\alpha_k} e^{-\lambda(t-t_j)} \\ &\quad \times \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\ &\quad + \sum_{j=1}^k \frac{(t-t_k)^{\alpha_k}}{\Gamma(\alpha_k+1)} \left[\frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\ &\quad + \frac{(t-t_k)^{\alpha_k} a_2}{\Gamma(\alpha_k+1)}, \end{aligned}$$

$$\begin{aligned} (18) \quad \sum_{k=0}^q \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} x(\eta_k) &= \sum_{k=0}^q \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} \left(\int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_{k-1}-1} h(s) ds \right) (\eta_k) + \sum_{k=0}^q \sum_{j=1}^k \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k-t_j)} \\ &\quad \times \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\ &\quad + \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k+1)} \\ &\quad \times \left[\frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\ &\quad + \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k} a_2}{\Gamma(\alpha_k+1)}. \end{aligned}$$

and

$$\begin{aligned} (19) \quad a_2 &= \left(1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k+1)} \right)^{-1} \\ &\quad \times \left\{ \sum_{k=0}^q \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} \left(\int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_{k-1}-1} h(s) ds \right) (\eta_k) \right. \\ &\quad + \sum_{k=0}^q \sum_{j=1}^k \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k-t_j)} \\ &\quad \times \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds - \frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\ &\quad \left. + \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k+1)} \left[\frac{1}{\lambda} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \right\}. \end{aligned}$$

Substituting the value of a_j ($j = 1, 2$) in (6) and (17), we obtain (5). Conversely, assume that x is a solution of the impulsive sequential fractional integral equation (5); then by a direct computation, it follows that the solution given by (5) satisfies (4). This completes the proof. \square

We presented some estimations that used in the forthcoming theorems.

Theorem 2.1. *For any $h \in PC([0, T], \mathbb{R})$ with $\|h\| = \sup_{t \in [0, T]} |h(t)|$, we have*

(i)

$$\begin{aligned} \left| \mathbf{I}_{t_k^+}^{\alpha_k} \left(\int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds \right) (t) \right| &\leq \mathbf{I}_{t_k^+}^{\alpha_k} \left| \int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds \right| (t) \\ &\leq \frac{(\eta_k - t_k)^{\alpha_k + \beta_k - 1}}{\Gamma(\alpha_k) \Gamma(\beta_k)} \|h\|_{PC}. \end{aligned}$$

(ii)

$$\left| \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds \right| \leq \frac{(t - t_k)^{\beta_k - 1}}{\lambda \Gamma(\beta_k)} \left(1 - e^{-\lambda(t - t_k)} \right) \|h\|_{PC},$$

(iii)

$$\left| \int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds \right| \leq \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} \left(1 - e^{-\lambda(t_j - t_{j-1})} \right) \|h\|_{PC},$$

(iv)

$$\left| \int_{t_{j-1}}^{t_j} \frac{(t_j - s)^{\beta_{j-1}-2}}{\Gamma(\beta_{j-1} - 1)} h(s) ds \right| \leq \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\Gamma(\beta_{j-1})} \|h\|_{PC},$$

(v)

$$\left| \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} \mathbf{I}_{t_k^+}^{\beta_k + \alpha_k - 1} h(s) ds \right| \leq \frac{(\eta_k - t_k)^{\beta_k + \alpha_k - 1}}{\lambda \Gamma(\beta_k + \alpha_k)} \left(1 - e^{-\lambda(\eta_k - t_k^+)} \right) \|h\|_{PC}$$

(vi)

$$\left| \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} \frac{(\eta_k - s)^{\alpha_k-1}}{\Gamma(\alpha_k)} h(s) ds \right| \leq \frac{(\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \|h\|_{PC}.$$

Proof. Obviously

(i)

$$\begin{aligned}
& \left| \mathbf{I}_{t_k^+}^{\alpha_k} \left(\int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds \right) (\eta_k) \right| \leq \mathbf{I}_{t_k^+}^{\alpha_k} \left| \int_{t_k}^r e^{-\lambda(r-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds \right| (\eta_k) \\
& \leq \int_{t_k}^{\eta_k} \frac{(\eta_k - r)^{\alpha_k-1}}{\Gamma(\alpha_k)} \\
& \quad \times \left(\left| \int_{t_k}^r e^{-\lambda(r-s)} \left(\int_{t_k}^s \frac{(s-u)^{\beta_k-2}}{\Gamma(\beta_k-1)} h(u) du \right) ds \right| \right) dr \\
& \leq \frac{(\eta_k - t_k)^{\beta_k-1}}{\Gamma(\beta_k)} \left(\int_{t_k}^{\eta_k} \frac{(\eta_k - r)^{\alpha_k-1}}{\Gamma(\alpha_k)} e^{-\lambda(\eta_k-r)} dr \right) \|h\|_{PC} \\
& \leq \frac{(\eta_k - t_k)^{\alpha_k-1}}{\Gamma(\alpha_k)} \frac{(\eta_k - t_k)^{\beta_k-1}}{\Gamma(\beta_k)} \left(\int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-r)} dr \right) \|h\|_{PC} \\
& \leq \frac{(\eta_k - t_k)^{\alpha_k+\beta_k-1}}{\Gamma(\alpha_k)\Gamma(\beta_k)} \|h\|_{PC}.
\end{aligned}$$

$$\begin{aligned}
(ii) \quad & \left| \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} h(s) ds \right| = \left| \int_{t_k}^t e^{-\lambda(t-s)} \left(\int_{t_k}^s \frac{(s-u)^{\beta_k-2}}{\Gamma(\beta_k-1)} h(u) du \right) ds \right| \\
& \leq \frac{(t - t_k)^{\beta_k-1}}{\Gamma(\beta_k)} \left(\int_{t_k}^t e^{-\lambda(t-s)} ds \right) \|h\|_{PC} \\
& \leq \frac{(t - t_k)^{\beta_k-1}}{\lambda \Gamma(\beta_k)} \left(1 - e^{-\lambda(t-t_k)} \right) \|h\|_{PC}, h \in PC(J, \mathbb{R}).
\end{aligned}$$

(iii)

$$\begin{aligned}
& \left| \int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(s) ds \right| = \left| \int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \left(\int_{t_{j-1}}^s \frac{(s-u)^{\beta_{j-1}-2}}{\Gamma(\beta_{j-1}-1)} h(u) du \right) ds \right| \\
& \leq \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\Gamma(\beta_k)} \left(\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} ds \right) \|h\|_{PC} \\
& \leq \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} \left(1 - e^{-\lambda(t_j-t_{j-1})} \right) \|h\|_{PC}, h \in PC(J, \mathbb{R}).
\end{aligned}$$

$$(iv) \quad \left| \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} h(t_j) \right| = \left| \int_{t_{j-1}}^{t_j} \frac{(t_j - s)^{\beta_{j-1}-2}}{\Gamma(\beta_{j-1}-1)} h(s) ds \right| \leq \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\Gamma(\beta_{j-1})} \|h\|_{PC}, h \in PC(J, \mathbb{R}).$$

(v)

$$\begin{aligned}
\left| \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} I_{t_k^+}^{\beta_k+\alpha_k-1} h(s) ds \right| &= \left| \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} \left(\int_{t_k}^s \frac{(s-u)^{\beta_k+\alpha_k-2}}{\Gamma(\beta_k+\alpha_k-1)} h(u) du \right) ds \right| \\
&\leq \frac{(\eta_k-t_k)^{\beta_k+\alpha_k-1}}{\Gamma(\beta_k+\alpha_k)} \left(\int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} ds \right) \|h\|_{PC} \\
&\leq \frac{(\eta_k-t_k)^{\beta_k+\alpha_k-1}}{\lambda \Gamma(\beta_k+\alpha_k)} \left(1 - e^{-\lambda(\eta_k-t_k)} \right) \|h\|_{PC}, h \in PC(J, \mathbb{R}).
\end{aligned}$$

(vi)

$$\begin{aligned}
\left| I_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k-t_j)} \right| &= \left| \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} \frac{(\eta_k-s)^{\alpha_k-1}}{\Gamma(\alpha_k)} h(s) ds \right| \\
&\leq \frac{(\eta_k-t_k)^{\alpha_k}}{\Gamma(\alpha_k+1)} \|h\|_{PC}, h \in PC(J, \mathbb{R}).
\end{aligned}$$

The proofs is completed. \square

3. MAIN RESULTS

This section deals with the existence and uniqueness of solutions for the problem (1). Before stating and proving the main results, we introduce the following hypotheses.

(H₁): there exist a nonnegative function $a(t) \in L(0, T)$ such that

$$|f(t, x)| \leq a(t) + \xi |x|^\sigma, \sigma > 0,$$

where ζ are nonnegative constant.

(H₂): there exists a constants L_ψ and L_{ψ^*} such that

$$|\psi_k(x)| \leq L_\psi, |\psi_k^*(x)| \leq L_{\psi^*}, t \in J, x \in \mathbb{R}, k = 1, 2, \dots, q.$$

(H₃): the function $f : J \times \mathbb{R} \rightarrow \mathbb{R}$ is jointly continuous

(H₄): there exists a constant $L_f > 0$ such that

$$|f(t, x) - f(t, y)| \leq L_f |x - y|, t \in J, x, y \in \mathbb{R}.$$

(H₅): there exist a positive constants L_ψ, L_{ψ^*} such that $|\psi_k(x) - \psi_k(y)| \leq L_\psi |x - y|, |\psi_k^*(x) - \psi_k^*(y)| \leq L_{\psi^*} |x - y|$.

Define an operator $\mathcal{F} : PC(J) \rightarrow PC(J)$ by

$$\begin{aligned}
(20) \quad \mathcal{F}x(t) = & \int_{t_k}^t e^{-\lambda(t-s)} I_{t_k^+}^{\beta_k-1} f(s, x(s)) ds \\
& + \sum_{j=1}^k e^{-\lambda(t-t_j)} \times \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_i-s)} I_{t_{j-1}^+}^{\beta_{j-1}-1} f(s, x(s)) ds \right. \\
& - \frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} f(t_j, x(t_j)) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \Big] \\
& + \sum_{j=1}^k \left[\frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} f(t_j, x(t_j)) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \\
& + \left(1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right)^{-1} \\
& \times \left\{ \sum_{k=0}^q \lambda_k I_{t_k^+}^{\alpha_k} \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} I_{t_k^+}^{\beta_k-1} f(s, x(s)) ds \right. \\
& + \sum_{k=0}^q \sum_{j=1}^k \lambda_k I_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k-t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} I_{t_{j-1}^+}^{\beta_{j-1}-1} f(s, x(s)) ds \right. \\
& - \frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} f(t_j, x(t_j)) - \frac{1}{\lambda} \psi_j^*(x(t_j)) \Big] \\
& \left. + \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\frac{1}{\lambda} I_{t_{j-1}^+}^{\beta_{j-1}-1} f(t_j, x(t_j)) + \psi_j(x(t_j)) + \frac{1}{\lambda} \psi_j^*(x(t_j)) \right] \right\},
\end{aligned}$$

For convenience, we will give some notations:

$$\begin{aligned}
\Delta &= \left| 1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right|^{-1}, \\
\Lambda_1 &= \sum_{j=1}^{q+1} \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})}, \quad \Lambda_2 = \sum_{j=1}^q \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} \Lambda_3 = \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k + \beta_k - 1}}{\Gamma(\alpha_k) \Gamma(\beta_k)}, \\
\Lambda_4 &= \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k} (t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\alpha_k + 1) \Gamma(\beta_{j-1})}, \quad \Lambda_5 = \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)}, \\
\mu(u) &= \Psi \|u\| + (1 + \Delta \Lambda_5) q L_\psi + \left(\frac{T}{\lambda} + \frac{1}{\lambda} + \left(\frac{1}{\lambda} + \frac{1}{\lambda} \right) \Delta \Lambda_5 \right) q L_{\psi^*}.
\end{aligned}$$

Theorem 3.1. Suppose that (H₃), (H₄) and (H₅) hold. If

$$(21) \quad L_{\mathcal{F}} < 1$$

then the equation (1) has a unique solution on J .

Proof. Show that $\mathcal{F} : PC(J) \rightarrow PC(J)$ is a completely continuous operator

$$\begin{aligned}
|\mathcal{F}x(t)| &\leq \int_{t_k}^t e^{-\lambda(t-s)} I_{t_k^+}^{\beta_k-1} |f(s, x(s))| ds \\
&+ \sum_{j=1}^k e^{-\lambda(t-t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_i-s)} I_{t_{j-1}^+}^{\beta_{j-1}-1} |f(s, x(s))| ds \right. \\
&+ \left. \left| \frac{1}{\lambda} \left| I_{t_{j-1}^+}^{\beta_{j-1}-1} |f(t_j, x(t_j))| + \left| \frac{1}{\lambda} \right| |\psi_j^*(x(t_j))| \right| \right] \\
&+ \sum_{j=1}^k \left[\left| \frac{1}{\lambda} \left| I_{t_{j-1}^+}^{\beta_{j-1}-1} |f(t_j, x(t_j))| + |\psi_j(x(t_j))| + \left| \frac{1}{\lambda} \right| |\psi_j^*(x(t_j))| \right| \right] \\
&+ \left| 1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right|^{-1} \\
&\times \left\{ \sum_{k=0}^q \lambda_k I_{t_k^+}^{\alpha_k} \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} I_{t_k^+}^{\beta_k-1} |f(s, x(s))| ds + \sum_{k=0}^q \sum_{j=1}^k \lambda_k I_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k-t_j)} \right. \\
&\times \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} I_{t_{j-1}^+}^{\beta_{j-1}-1} |f(s, x(s))| ds + \left| \frac{1}{\lambda} \left| I_{t_{j-1}^+}^{\beta_{j-1}-1} |f(t_j, x(t_j))| + \left| \frac{1}{\lambda} \right| |\psi_j^*(x(t_j))| \right| \right] \\
&+ \left. \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\left| \frac{1}{\lambda} \left| I_{t_{j-1}^+}^{\beta_{j-1}-1} |f(t_j, x(t_j))| + |\psi_j(x(t_j))| + \left| \frac{1}{\lambda} \right| |\psi_j^*(x(t_j))| \right| \right] \right\},
\end{aligned}$$

$$\begin{aligned}
|\mathcal{F}x(t)| &\leq L_f \frac{(t-t_k)^{\beta_k-1}}{\lambda \Gamma(\beta_k)} \\
&+ \sum_{j=1}^k e^{-\lambda(t-t_j)} \left[L_f \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_f \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_{\psi^*} \frac{1}{\lambda} \right] \\
&+ \sum_{j=1}^k \left[L_f \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_{\psi} + L_{\psi^*} \frac{1}{\lambda} \right] + \Delta \left\{ L_f \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k + \beta_k - 1}}{\Gamma(\alpha_k) \Gamma(\beta_k)} \cdot \right. \\
&+ \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[L_f \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_f \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_{\psi^*} \frac{1}{\lambda} \right] \\
&+ \left. \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[L_f \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_{\psi} + L_{\psi^*} \frac{1}{\lambda} \right] \right\},
\end{aligned}$$

$$\begin{aligned}
|\mathcal{F}x(t)| &\leq L_f \frac{(t-t_k)^{\beta_k-1}}{\lambda \Gamma(\beta_k)} + L_f \sum_{j=1}^k e^{-\lambda(t-t_j)} \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} \\
&+ L_f \sum_{j=1}^k e^{-\lambda(t-t_j)} \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + \frac{1}{\lambda} \sum_{j=1}^k e^{-\lambda(t-t_j)} L_{\psi^*} \\
&+ L_f \sum_{j=1}^k \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + \sum_{j=1}^k L_\psi + \frac{1}{\lambda} \sum_{j=1}^k L_{\psi^*} + \Delta \left\{ L_f \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k + \beta_k - 1}}{\Gamma(\alpha_k) \Gamma(\beta_k)} \right. \\
&+ \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} L_f + L_f \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + \frac{1}{\lambda} \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} L_{\psi^*} \right] \\
&+ \left. \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} L_f + L_\psi + \frac{1}{\lambda} L_{\psi^*} \right] \right\},
\end{aligned}$$

$$\begin{aligned}
|\mathcal{F}x(t)| &\leq L_f \sum_{j=1}^{q+1} \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + T L_f \sum_{j=1}^q \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + T L_f \sum_{j=1}^q \frac{(t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} \\
&+ \frac{qT}{\lambda} L_{\psi^*} + q L_\psi + \frac{q}{\lambda} L_{\psi^*} + \Delta \left\{ L_f \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k + \beta_k - 1}}{\Gamma(\alpha_k) \Gamma(\beta_k)} \right. \\
&+ L_f \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k} (t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\alpha_k + 1) \Gamma(\beta_{j-1})} + L_f \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k} (t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\alpha_k + 1) \Gamma(\beta_{j-1})} \\
&+ \frac{1}{\lambda} \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} L_{\psi^*} + L_f \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k} (t_j - t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\alpha_k + 1) \Gamma(\beta_{j-1})} \\
&+ \left. \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} L_\psi + \frac{1}{\lambda} \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} L_{\psi^*} \right\}, \\
&\leq L_f \{(1+q)\Lambda_1 + qT\Lambda_2 + qT\Lambda_2\} + \frac{qT}{\lambda} L_{\psi^*} + q L_\psi + \frac{q}{\lambda} L_{\psi^*} \\
&+ \Delta \left\{ L_f \Lambda_3 + q L_f \Lambda_4 + q L_f \Lambda_4 + \frac{q}{\lambda} \Lambda_5 L_{\psi^*} + q L_f \Lambda_4 + q \Lambda_5 L_\psi + \frac{q}{\lambda} \Lambda_5 L_{\psi^*} \right\}, \\
&\leq L_f \{((1+q)\Lambda_1 + 2qT\Lambda_2) + \Delta(\Lambda_3 + 3q\Lambda_4)\} \\
&+ (1 + \Delta\Lambda_5) q L_\psi + \left(\frac{T}{\lambda} + \frac{1}{\lambda} + \left(\frac{1}{\lambda} + \frac{1}{\lambda} \right) \Delta\Lambda_5 \right) q L_{\psi^*}
\end{aligned}$$

which implies

$$|\mathcal{F}x(t)| \leq \Psi L_f + (1 + \Delta\Lambda_5) qL_\psi + \left(\frac{T}{\lambda} + \frac{1}{\lambda} + \left(\frac{1}{\lambda} + \frac{1}{\lambda} \right) \Delta\Lambda_5 \right) qL_{\psi^*}.$$

which implies that $\mathcal{F}x \in B$. Thus $\mathcal{F}B \subset B$. On the other hand, for any $t \in J_k, 0 \leq k \leq q$, we have

$$\begin{aligned} |(\mathcal{F}x)'(t)| &\leq \lambda \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} |f(s, x(s))| ds + \int_{t_k}^t \frac{(t-s)^{\beta_k-2}}{\Gamma(\beta_k-1)} |f(s, x(s))| ds \\ &+ \sum_{j=1}^k \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} |f(s, x(s))| ds \right. \\ &\quad \left. + \frac{1}{\lambda} \int_{t_{j-1}}^{t_j} \frac{(t_j-s)^{\beta_{j-1}-2}}{\Gamma(\beta_{j-1}-1)} |f(s, x(s))| ds + \left| \frac{1}{\lambda} \psi_j^*(x(t_j)) \right| \right], \\ &\leq \lambda L_f \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k^+}^{\beta_k-1} ds + L_f \int_{t_k}^t \frac{(t-s)^{\beta_k-2}}{\Gamma(\beta_k-1)} ds \\ &+ \sum_{j=1}^q \left[L_f \int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1}-1} ds + L_f \frac{1}{\lambda} \int_{t_{j-1}}^{t_j} \frac{(t_j-s)^{\beta_{j-1}-2}}{\Gamma(\beta_{j-1}-1)} ds + \frac{1}{\lambda} L_{\psi^*} \right], \\ &\leq L_f \left[\frac{(t-t_k)^{\beta_k-1}}{\Gamma(\beta_k)} \left((1 - e^{-\lambda(t-t_k)}) + 1 \right) \right] \\ &+ \frac{q}{\lambda} \left[L_f \left[\frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\Gamma(\beta_k)} \left((1 - e^{-\lambda(t_j-t_{j-1})}) + 1 \right) \right] + L_{\psi^*} \right] \\ &= \mathcal{L}. \end{aligned}$$

Hence, for $\tau_1, \tau_2 \in J_k$ with $\tau_1 \leq \tau_2$ and $0 \leq k \leq q$, we have

$$|(\mathcal{F}x)(\tau_2) - (\mathcal{F}x)(\tau_1)| \leq \int_{\tau_1}^{\tau_2} |(\mathcal{F}x)'(s)| ds \leq \mathcal{L}(\tau_2 - \tau_1).$$

This implies that $\mathcal{F}x$ is equicontinuous on all $J_k, k = 0, 1, \dots, q$. Consequently, Arzela-Ascoli theorem ensures the operator $\mathcal{F} : PC(J, \mathbb{R}) \rightarrow PC(J, \mathbb{R})$ is a completely continuous operator. Next show that the operator \mathcal{F} maps B into B . For that, let us choose $R \geq \max \left\{ 2\mu, (2L_\sigma)^{\frac{1}{1-\sigma}} \right\}$ and define a ball $B = \{x \in PC(J, \mathbb{R}) : \|x\| \leq R\}$. For any $x \in B$, by the conditions (H_1) and

(H_2) , we have

$$\begin{aligned}
|\mathcal{F}x(t)| &\leq \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k}^{\beta_k-1} [a(s) + \xi |x(s)|^\sigma] ds \\
&+ \sum_{j=1}^k e^{-\lambda(t-t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [a(s) + \xi |x(s)|^\sigma] ds \right. \\
&+ \left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [a(t_j) + \xi |x(t_j)|^\sigma] \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) \right| \right| \right] \\
&+ \sum_{j=1}^k \left[\left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [a(t_j) + \xi |x(t_j)|^\sigma] \right| + \left| \psi_j(x(t_j)) \right| \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) \right| \right] \\
&+ \left| 1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right|^{-1} \times \left\{ \sum_{k=0}^q \lambda_k \mathbf{I}_{t_k}^{\alpha_k} \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k-s)} \mathbf{I}_{t_k}^{\beta_k-1} [a(s) + \xi |x(s)|^\sigma] ds \right. \\
&+ \sum_{k=0}^q \sum_{j=1}^k \lambda_k \mathbf{I}_{t_k}^{\alpha_k} e^{-\lambda(\eta_k-t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [a(s) + \xi |x(s)|^\sigma] ds \right. \\
&+ \left. \left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [a(t_j) + \xi |x(t_j)|^\sigma] \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) \right| \right| \right] \\
&+ \left. \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [a(t_j) + \xi |x(t_j)|^\sigma] \right| + \left| \psi_j(x(t_j)) \right| \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) \right| \right] \right\},
\end{aligned}$$

$$\begin{aligned}
|\mathcal{F}x(t)| &\leq [\|a\| + \xi \|x\|^\sigma] \frac{(t-t_k)^{\beta_k-1}}{\lambda \Gamma(\beta_k)} + \sum_{j=1}^k e^{-\lambda(t-t_j)} \\
&\times \left[[\|a\| + \xi \|x\|^\sigma] \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + [\|a\| + \xi \|x\|^\sigma] \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + \frac{1}{\lambda} L_{\psi^*} \right] \\
&+ \sum_{j=1}^k \left[[\|a\| + \xi \|x\|^\sigma] \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_\psi + \frac{1}{\lambda} L_{\psi^*} \right] \\
&+ \Delta \left\{ [\|a\| + \xi \|x\|^\sigma] \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k + \beta_k - 1}}{\Gamma(\alpha_k) \Gamma(\beta_k)} + \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right. \\
&\times \left[[\|a\| + \xi \|x\|^\sigma] \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + [\|a\| + \xi \|x\|^\sigma] \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + \frac{1}{\lambda} L_{\psi^*} \right] \\
&+ \left. \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[[\|a\| + \xi \|x\|^\sigma] \frac{(t_j-t_{j-1})^{\beta_{j-1}-1}}{\lambda \Gamma(\beta_{j-1})} + L_\psi + \frac{1}{\lambda} L_{\psi^*} \right] \right\},
\end{aligned}$$

$$\begin{aligned} &\leq \Psi \|a\| + (1 + \Delta\Lambda_5) qL_\psi + \left(\frac{T}{\lambda} + \frac{1}{\lambda} + \left(\frac{1}{\lambda} + \frac{1}{\lambda} \right) \Delta\Lambda_5 \right) qL_{\psi^*} + \Psi\xi \|x\|^\sigma. \\ &\leq \mu(a) + \Psi\xi \|x\|^\sigma. \end{aligned}$$

Thus,

$$|\mathcal{F}x(t)| \leq \mu(a) + \Psi\xi \|x\|^\sigma \leq \frac{R}{2} + \frac{R}{2} = R.$$

This implies $\mathcal{F} : B \rightarrow B$. Hence, we conclude that $\mathcal{F} : B \rightarrow B$ is completely continuous. It follows from the Schauder fixed point theorem that the operator \mathcal{F} has at least one fixed point. That is problem (1) has at least one solution in B .

Theorem 3.2. Assume that there exist a nonnegative function $W \in C(J, \mathbb{R}^+)$ and nonnegative constants M, Z such that

$$|f(t, x) - f(t, y)| \leq W(t) |x - y|, \quad t \in J, \quad x, y \in \mathbb{R},$$

$$|\psi_k(x) - \psi_k(y)| \leq M|x - y|, \quad |\psi_k^*(x) - \psi_k^*(y)| \leq Z|x - y|,$$

for $t \in J, x, y \in \mathbb{R}$ and $k = 1, 2, \dots, q$. Furthermore, the assumption $\mu(W) < 1$ holds. Then the equation (1) has a unique solution on J .

□

Proof. For $x, y \in B$ and for each $t \in J$, we have

$$\begin{aligned} |(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| &\leq \int_{t_k}^t e^{-\lambda(t-s)} \mathbf{I}_{t_k}^{\beta_k-1} |f(s, x(s)) - f(s, y(s))| ds \\ &\quad + \sum_{j=1}^k e^{-\lambda(t-t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j-s)} \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} |f(s, x(s)) - f(s, y(s))| ds \right. \\ &\quad \left. + \left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [f(t_j, x(t_j)) - f(t_j, y(t_j))] \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) - \psi_j^*(y(t_j)) \right| \right| \right] \\ &\quad + \sum_{j=1}^k \left[\left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}}^{\beta_{j-1}-1} [f(t_j, x(t_j)) - f(t_j, y(t_j))] \right| + \left| \psi_j(x(t_j)) - \psi_j(y(t_j)) \right| \right| \right. \\ &\quad \left. + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) - \psi_j^*(y(t_j)) \right| \right| \right] \end{aligned}$$

$$\begin{aligned}
& + \left| 1 - \sum_{k=0}^q \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \right|^{-1} \left\{ \sum_{k=0}^q \lambda_k \mathbf{I}_{t_k}^{\alpha_k} \int_{t_k}^{\eta_k} e^{-\lambda(\eta_k - s)} \mathbf{I}_{t_k}^{\beta_k - 1} |f(s, x(s)) - f(s, y(s))| ds \right. \\
& + \sum_{k=0}^q \sum_{j=1}^k \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} e^{-\lambda(\eta_k - t_j)} \left[\int_{t_{j-1}}^{t_j} e^{-\lambda(t_j - s)} \mathbf{I}_{t_{j-1}}^{\beta_{j-1} - 1} |f(s, x(s)) - f(s, y(s))| ds \right. \\
& + \left. \left. + \left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1} - 1} |f(t_j, x(t_j)) - f(t_j, y(t_j))| \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) - \psi_j^*(y(t_j)) \right| \right| \right] \right\} \\
& + \sum_{k=0}^q \sum_{j=1}^k \frac{\lambda_k (\eta_k - t_k)^{\alpha_k}}{\Gamma(\alpha_k + 1)} \left[\left| \frac{1}{\lambda} \left| \mathbf{I}_{t_{j-1}^+}^{\beta_{j-1} - 1} |f(t_j, x(t_j)) - f(t_j, y(t_j))| \right| \right. \right. \\
& + \left. \left. + \left| \psi_j(x(t_j)) - \psi_j(y(t_j)) \right| + \left| \frac{1}{\lambda} \left| \psi_j^*(x(t_j)) - \psi_j^*(y(t_j)) \right| \right| \right] \right\}, \\
& \leq \left\{ \Psi \|W\| + (1 + \Delta\Lambda_5) q L_\psi + \left(\frac{T}{\lambda} + \frac{1}{\lambda} + \left(\frac{1}{\lambda} + \frac{1}{\lambda} \right) \Delta\Lambda_5 \right) q L_{\psi^*} \right\} \|x - y\|. \\
& = \mu(W) \|x - y\|.
\end{aligned}$$

As $\mu(W) \leq 1$, we have $|(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| < \|x - y\|$. Therefore, \mathcal{F} is a contraction mapping on $PC(J, \mathbb{R})$ due to condition (21). By applying the well-known Banach's contraction mapping we see that the operator \mathcal{F} has a unique fixed point on $PC(J, \mathbb{R})$. Therefore, the problem (1) has a unique solution. This completes the proof.

Example (1) consider the impulsive sequential fractional deferential equation

□

$$\begin{aligned}
(22) \quad & \left({}^c D_{t_k^+}^{\beta_k} + \lambda {}^c D_{t_k^+}^{\beta_k - 1} \right) x(t) = \frac{e^t \sin [3x(t) + e^{(\frac{1}{2})x(t)}]}{2 + x^4(t)} + \frac{\cos(2t + 5)}{\sqrt{3 + x^2(t)}} |x(t)|^\sigma, \\
& 0 < t \leq 1, t \neq \frac{3}{4}, k = 0, 1, \dots, q; \\
& \Delta x\left(\frac{3}{4}\right) = 11 \sin^2 x\left(\frac{1}{4}\right), \Delta x'\left(\frac{3}{4}\right) = \frac{|x\left(\frac{3}{4}\right)|}{2(1 + |x\left(\frac{3}{4}\right)|)}, \\
& x(0) = \sum_{k=0}^1 \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} x(\eta_k) + \frac{1}{2}, x'(0) = 0.
\end{aligned}$$

$t \in [0, 1]$, let $\beta_0 = \frac{5}{4}, \beta_1 = 1, \beta = \frac{8}{5}, \alpha_0 = \frac{1}{2}, \alpha_1 = \frac{5}{3}, \lambda_0 = \frac{2}{5}, \lambda_1 = \frac{3}{7}, \eta_1 = \frac{1}{2}, \eta_2 = \frac{4}{5}$. Observe that

$$\begin{aligned} |(t, x, y)| &= \left| \frac{e^t \sin [3x(t) + e^{(\frac{1}{2})x(t)}]}{2+x^4(t)} + \frac{\cos(2t+5)}{\sqrt{3+x^2(t)}} |x(t)|^\sigma \right| \\ &\leq \frac{e^t}{2} + \frac{1}{\sqrt{3}} |x| \sigma. \end{aligned}$$

Clearly, $a(t) = \frac{e^t}{2}$, $\xi = \frac{1}{\sqrt{3}}$, $L_\psi = 11$, $L_{\psi^*} = \frac{1}{2}$, and the conditions of Theorem 4 hold. Thus, by Theorem, the impulsive sequential multi-orders fractional boundary value problem (22) has at least one solution.

Example (2) consider the impulsive sequential fractional deferential equation

$$\begin{aligned} ({}^cD_{t_k^+}^{\beta_k} + \lambda {}^cD_{t_k^+}^{\beta_k-1})x(t) &= \frac{e^t \sin [2x(t) + e^{(\frac{1}{2})x(t)}]}{2+x^4(t)} + \left[\left(\frac{\cos(2t+5)}{\sqrt{1+x^2(t)}} * \frac{\arctan^2 x(t)}{\sqrt{2+x(t)}} \right) |x(t)|^\sigma \right], \\ (22) \quad 0 < t \leq 1, t \neq \frac{1}{3}, k &= 0, 1, \dots, q; \\ \Delta x(\frac{1}{3}) &= 11 \sin^2 x\left(\frac{1}{3}\right), \Delta x'(\frac{1}{3}) = \frac{|x(\frac{1}{3})|}{2(1+|x(\frac{1}{3})|)}, \\ x(0) &= \sum_{k=0}^1 \lambda_k \mathbf{I}_{t_k^+}^{\alpha_k} x(\eta_k) + \frac{1}{2}, x'(0) = 0. \end{aligned}$$

$t \in [0, 1]$, let $\beta_0 = \frac{3}{4}$, $\beta_1 = 1$, $\beta = \frac{6}{5}$, $\alpha_0 = \frac{1}{2}$, $\alpha_1 = \frac{2}{3}$, $\lambda_0 = \frac{1}{5}$, $\lambda_1 = \frac{3}{5}$, $\eta_1 = \frac{1}{2}$, $\eta_2 = \frac{4}{5}$.

Observe that

$$\begin{aligned} |(t, x, y)| &= \frac{e^t \sin [2x(t)]}{2+x^2(t)} + \left[\left(\frac{\cos(2t+4)}{\sqrt{1+x(t)}} * \frac{\arctan^2 x(t)}{\sqrt{2+x(t)}} \right) |x(t)|^\sigma \right] \\ &\leq \frac{e^t}{2} + \frac{4}{\sqrt{3}} |x| \sigma. \end{aligned}$$

CONCLUSION

In this paper discussed a new impulsive sequential multi-orders fractional differential equation with boundary condition. The existence and uniqueness result are obtaind for a nonlinear proplem with fractionL integral boundary condition applying standard fixed point theorems. So the present work is a useful contribution to the existing literature on the topic.

CONFLICT OF INTERESTS

The author(s) declare that there is no conflict of interests.

REFERENCES

- [1] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and Applications of Fractional Differential Equations. North-Holland Mathematics Studies, vol. 204. Elsevier, Amsterdam, 2006.
- [2] V. Lakshmikantham, S. Leela, D.J. Vasundhara, Theory of Fractional Dynamic Systems, Cambridge Scientific Publisher, Cambridge, 2009.
- [3] J. Sabatier, O.P. Agrawal, J.A.T. Machado, Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering. Springer, Dordrecht; 2007.
- [4] D. Baleanu, K. Diethelm, E. Scalas, J.J. Trujillo, Fractional Calculus Models and Numerical Methods, Series on Complexity, Nonlinearity and Chaos, World Scientific, Singapore, 2012.
- [5] E. Dinç, D. Baleanu, Fractional wavelet transform for the quantitative spectral resolution of the composite signals of the active compounds in a two-component mixture, *Comput. Math. Appl.* 59 (2010) 1701–1708.
- [6] G. Wang, S. Liu, D. Baleanu, L. Zhang, A New Impulsive Multi-Orders Fractional Differential Equation Involving Multipoint Fractional Integral Boundary Conditions, *Abstr. Appl. Anal.* 2014 (2014), 932747.
- [7] B. Ahmad, S. Sivasundaram, Existence results for nonlinear impulsive hybrid boundary value problems involving fractional differential equations, *Nonlinear Anal., Hybrid Syst.* 3 (2009), 251–258.
- [8] A. Debbouche, D. Baleanu, Controllability of fractional evolution nonlocal impulsive quasilinear delay integro-differential systems, *Comput. Math. Appl.* 62 (2011), 1442–1450.
- [9] Y. Tian, Z. Bai, Existence results for the three-point impulsive boundary value problem involving fractional differential equations, *Comput. Math. Appl.* 59 (2010), 2601–2609.
- [10] G.M. Mophou, Existence and uniqueness of mild solutions to impulsive fractional differential equations, *Nonlinear Anal., Theory Meth. Appl.* 72 (2010), 1604–1615.
- [11] C. Bai, Impulsive periodic boundary value problems for fractional differential equation involving Riemann–Liouville sequential fractional derivative, *J. Math. Anal. Appl.* 384 (2011), 211–231.
- [12] L. Zhang, G. Wang, G. Song, Existence of Solutions for Nonlinear Impulsive Fractional Differential Equations of Order $\alpha \in (2, 3]$ with Nonlocal Boundary Conditions, *Abstr. Appl. Anal.* 2012 (2012), 717235.
- [13] L. Zhang and G. Wang, Existence of solutions for nonlinear fractional differential equations with impulses and anti-periodic boundary conditions, *Electron. J. Qual. Theory Differ. Equ.* 2011 (2011), 7.
- [14] Z. Liu, L. Lu, I. Szántó, Existence of solutions for fractional impulsive differential equations with p-Laplacian operator, *Acta Math. Hung.* 141 (2013), 203–219.

- [15] X. Zhang, X. Huang, Z. Liu, The existence and uniqueness of mild solutions for impulsive fractional equations with nonlocal conditions and infinite delay, *Nonlinear Anal., Hybrid Syst.* 4 (2010), 775–781.
- [16] M. A. Hammad, R. Khalil, Abel's formula and Wronskian for conformable fractional differential equations, *Int. J. Differ. Equ. Appl.* 13 (2014), 177–183.
- [17] K. Balachandran, S. Kiruthika, J.J. Trujillo, Existence results for fractional impulsive integrodifferential equations in Banach spaces, *Commun. Nonlinear Sci. Numer. Simul.* 16 (2011), 1970–1977.
- [18] B. Sami, On Impulsive Sequential Fractional Differential Equations with Separated Boundary Conditions, *Nonlinear Anal. Differ. Equ.* 5 2017, 283–297.
- [19] S. Kumar, A new efficient algorithm to solve non-linear fractional Ito coupled system and its approximate solution, *Walailak J. Sci. Technol.* 11 (2014), 1057–1067.
- [20] M. Khader, S. Kumar, S. Abbasbandy, Fractional homotopy analysis transforms method for solving a fractional heat-like physical model, *Walailak J. Sci. Technol.* 13 (2015), 337–353.