



Available online at <http://scik.org>

J. Math. Comput. Sci. 11 (2021), No. 6, 7082-7088

<https://doi.org/10.28919/jmcs/6182>

ISSN: 1927-5307

HYBRID ALGORITHM TO NEWTON RAPHSON METHOD AND BISECTION METHOD

CHALERMWUT COMEMUANG¹, PAIRAT JANNGAM^{2,*}

¹Department of Mathematics, Faculty of Science, Buriram Rajabhat University, Buriram, Thailand

²Department of Mathematics, Faculty of Education, Buriram Rajabhat University, Buriram, Thailand

Copyright © 2021 the author(s). This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract: The purpose of this research propose is to hybrid algorithm to Newton-Raphson method and Bisection method to compute roots of nonlinear equations. Numerical experiments for various tests nonlinear equations confirm performance for the bracketing method or open method to be compared with.

Keywords: Bisection method; Newton Raphson method; nonlinear equations.

2010 AMS Subject Classification: 65H05.

1. INTRODUCTION

Computing roots of nonlinear equations by iteration is a significant topic for numerical analysis. Because of the complicated problems of science, computer science and engineering, it cannot find the exact answer by the methods for solving equation. So, iteration is an alternative

*Corresponding author

E-mail address: pairat.jn@bru.ac.th

Received May 31, 2021

choice to find the answer or it could be called the step of finding root of function x which $f(x) = 0$ it is one of the well-known iteration method for finding answers called Newton-Raphson method and Bisection method. The derivative of function must be used for calculating the answer of every conducting iteration and has square convergence which is said to be the effective way to find the answer quickly. Then, there are a lot of researchers improving Newton-Raphson Method to be more convergent and one of them was Homeier[1] who presented the adjusted Newton-Raphson Method to have cubic convergence. Changbum and Beny[2] had improved fourth-order convergence. G. Mahesh[3] proposed a new method to compute a non-zero real root of the transcendental equations. Besides, there is a bisection method which is marginal specification with the notion of finding answers by using the intermediate value theorem to create closed interval containing the answers. The convergent sequence is linear, and it can certainly find the answer. However, this method could be used to find answer quite slowly, so Sabharwal[4] had improved bisection and false position method for finding the answers more swiftly. Kouider[5] presented a method of improved bisection and false position for many square root values through R program. Ali et al.[6] had improved bisection method by divided n as sub range. Tanakan[7] had improved bisection through the notion of Secant Method to find the answer rapidly. Neamvonk[8] has been a new method for solving nonlinear equations by using nonlinear regression. Hafiz [9] proposed an improved method called BRFC, by the combination of the Bisection, Regula Falsi and parabolic interpolation.

However, the method of marginal specification, square roots converged quite slowly, but Newton-Raphson method cannot find the certain answers depending on choosing appropriate beginning point, so Altaee, Hoomod and Hussein[10] presented hybrid algorithm among Newton-Raphson method. Kim, Noh, Oh and Park[11] had created the step of hybrid which used bisection and Newton-Raphson method to improve hybrid method of Altaee et al.

Therefore, this study revealed that the improvement of hybrid among Newton-Raphson method and bisection are more effective for convergence than interval or open approach.

2. MAIN RESULTS

Let f be a continuous function and defined on $[a, b]$ which $f(a) \cdot f(b) < 0$. Firstly, we set

$$a_1 = a - \frac{f(a)}{f'(a)}$$

$$b_1 = b - \frac{f(b)}{f'(b)}$$

we have two main cases:

case 1: If a_1 or $b_1 \notin [a, b]$, By the bisection method, we have $c = \frac{a+b}{2}$. Next, we consider a

new subinterval (a^*, b^*) by

$$(a^*, b^*) = \begin{cases} a^* = c, b^* = b & , \text{if } f(a) \cdot f(c) < 0 \\ a^* = a, b^* = c & , \text{if } f(a) \cdot f(c) > 0 \end{cases}$$

case 2: If a_1 or $b_1 \in [a, b]$

case 2.1: If $f(a_1) \cdot f(b_1) < 0$ then $c = \frac{a_1 + b_1}{2}$. Next, we consider a new subinterval

(a^*, b^*) by

$$(a^*, b^*) = \begin{cases} a^* = a_1, b^* = b & , \text{if } f(a_1) \cdot f(c) < 0 \\ a^* = c, b^* = b_1 & , \text{if } f(a_1) \cdot f(c) > 0 \end{cases}$$

case 2.2: If $f(a_1) \cdot f(b_1) > 0$, we set $a^* = a_1, b^* = b_1$ and

$$c = \begin{cases} a_1 & , \text{if } |f(a_1)| < |f(b_1)| \\ b_1 & , \text{if } |f(a_1)| > |f(b_1)| \end{cases}$$

Finally, we choose the new subinterval for the next iteration as follows $a = a^*$ and $b = b^*$

The process is continued until the interval is sufficiently small or the approximate solution is sufficiently close to the exact solution.

Therefore, we can state the algorithm for finding a solution of nonlinear equation $f(x) = 0$ on an interval $[a, b]$ as follows:

2.1. The algorithm

Hybrid Algorithm (HA)

Step 1: Get $f(x), f'(x), [a, b]$ (interval)

$$\text{Step 2: } a1 = a - \frac{f(a)}{f'(a)}, b1 = b - \frac{f(b)}{f'(b)}$$

Step 3: $a1$ or $b1 \notin [a, b]$, compute $c = \frac{a + b}{2}$ and

$$(a^*, b^*) = \begin{cases} a^* = c, b^* = b & , \text{if } f(a) \cdot f(c) < 0 \\ a^* = a, b^* = c & , \text{if } f(a) \cdot f(c) > 0 \end{cases}$$

go to Step 6.

Step 4: $f(a1) \cdot f(b1) < 0$ compute $c = \frac{a1 + b1}{2}$

$$(a^*, b^*) = \begin{cases} a^* = a1, b^* = c & , \text{if } f(a1) \cdot f(c) < 0 \\ a^* = c, b^* = b1 & , \text{if } f(a1) \cdot f(c) > 0 \end{cases}$$

go to Step 6.

Step 5: $f(a1) \cdot f(b1) > 0$, we set $a^* = a1, b^* = b1$ and

$$c = \begin{cases} a1 & , \text{if } |f(a1)| < |f(b1)| \\ b1 & , \text{if } |f(a1)| > |f(b1)| \end{cases}$$

Step 6: If $f(c) < \varepsilon$ then the zero is c . Stop.

Step 7: Set $a = a^*$ and $b = b^*$ and to Step 2.

3. NUMERICAL EXAMPLES

In this section, we compare our proposed methods Bisection Algorithm(BS)[12], Regula Falsi Algorithm(RF)[13], Improved Regula Falsi Algorithm(IRF)[13], Bisection Newton-like Algorithm(AC)[14], Newton-Raphson Algorithm(NR), Proposed Method(PP) and Hybrid Algorithm(HA) we tested the algorithms with the specific examples on Scilab (version 6.1.0)

program. For the accuracy, we use tolerance error(TOL) less than $\varepsilon = 1.0 \times 10^{-10}$ and also the

following criteria is used for estimating the zero $|f(x)| < \varepsilon$

$f(x)$	Methods	Interval	No. of iteration	$ f(x) $
$f_1(x) = x^2 - (1-x)^5$	BS	[0.1, 1]	33	3.29e-11
	RF	[0.1, 1]	20	4.83e-11
	IRF	[0.1, 1]	5	1.74e-14
	AC	[0.1, 1]	6	3.22e-13
	NR	0.1	5	0.00e+00
	PP	0.1	4	6.93e-17
	HA	[0.1, 1]	4	8.78e-12
$f_2(x) = \cos(x) - x^3$	BS	[0.1, 1]	34	4.65e-11
	RF	[0.1, 1]	12	3.59e-11
	IRF	[0.1, 1]	5	1.82e-11
	AC	[0.1, 1]	34	4.65e-11
	NR	0.1	10	2.95e-12
	PP	0.1	19	1.17e-12
	HA	[0.1, 1]	4	5.22e-15
$f_3(x) = xe^x - 1$	BS	[-1, 1]	34	2.84e-11
	RF	[-1, 1]	22	3.37e-11
	IRF	[-1, 1]	6	3.33e-16
	AC	[-1, 1]	8	2.27e-11
	NR	-1	2	0.00e+00
	PP	-1	1	0.00e+00
	HA	[-1, 1]	5	2.22e-16
$f_4(x) = x^2 - e^x - 3x + 2$	BS	[-2, 2]	36	4.95e-11
	RF	[-2, 2]	19	9.11e-11
	IRF	[-2, 2]	4	1.09e-13
	AC	[-2, 2]	36	4.95e-11
	NR	-2	5	0.00e+00
	PP	-2	∞	∞
	HA	[-2, 2]	3	1.15e-12

HYBRID ALGORITHM METHOD

$f(x)$	Methods	Interval	No. of iteration	$ f(x) $
$f_5(x) = \sin(x) - \frac{x}{2}$	BS	[-3, 5]	35	6.36e-11
	RF	[-3, 5]	24	4.32e-11
	IRF	[-3, 5]	11	1.65e-11
	AC	[-3, 5]	35	6.36e-11
	NR	-3	5	0.00e+00
	PP	-3	5	1.32e-13
	HA	[-3, 5]	5	0.00e+00
$f_6(x) = x^3$	BS	[-0.5, 1/3]	9	3.44e-11
	RF	[-0.5, 1/3]	579117	1.00e-10
	IRF	[-0.5, 1/3]	463465	1.00e-10
	AC	[-0.5, 1/3]	7	7.84e-11
	NR	-0.5	18	3.87e-11
	PP	-0.5	20	3.93e-11
	HA	[-0.5, 1/3]	7	4.43e-13

4. CONCLUSIONS

We presented a class of Hybrid Algorithm methods for finding simple zeros of nonlinear equations. In this paper, an algorithm Hybrid Algorithm is developed for computational purposes. The algorithm is tested on a number of numerical examples and the results obtain up to the desire accuracy are compared with the other methods. It is observed that our method takes a smaller number of iterations and more effective in comparison with these methods.

5. ACKNOWLEDGMENT

The authors wish to thank the referees for their interesting suggestions and comments to improve this article.

6. CONFLICT OF INTERESTS

The authors declare that there is no conflict of interests.

REFERENCES

- [1] H. Homeier, A modified newton method for root finding with cubic convergence, *J. Comput. Appl. Math.* 157 (2003), 227-230.
- [2] C. Changbum, N. Beny, Certain improvements of newtons method with fourth-order convergence, *Appl. Math. Comput.* 215 (2009), 821-828.
- [3] G. Mahesh, G. Swapna, K. Venkateshwarlu, An iterative method for solving non-linear transcendental equations, *J. Math. Comput. Sci.* 10 (2020), 1633-1642.
- [4] C. L. Sabharwal, Blended root finding algorithm outperforms bisection and regula falsi algorithms, *Mathematics.* 2019 (2019), 1118.
- [5] M. R. Kouider, Modified bisection algorithm for multiple roots of nonlinear equation with the R software, *SSRN.* (2019), <https://dx.doi.org/10.2139/ssrn.3451155>.
- [6] M. R. M. Ali, M. I. Fakhri, N. Hayati, N. A. Ramli, I. Jusoh, The n-th section method: A modification of bisection, *Malays. J. Fundam. Appl. Sci.* 13 (2017), 728–731.
- [7] S. Tanakan, A new algorithm of modified bisection method for nonlinear equation, *Appl. Math. Sci.* 123 (2013), 6107-6114.
- [8] J. Neamvonk, B. Phuenaree, A. Neamvonk, A new method for finding root of nonlinear equations by using nonlinear regression, *Asian J. Appl. Sci.* 6 (2015), 818-822.
- [9] M. Hafiz, A new combined bracketing method for solving nonlinear equations, *J. Math. Comput. Sci.* 3 (2013), 87–93.
- [10] A. A. H. Altaee, H. K. Hoomod, K. A. Hussein, A new approach to find roots of nonlinear equations by hybrid algorithm to bisection and Newton-Raphson algorithms, *Iraq. J. Inform. Technol.* 7 (2015), 75–82.
- [11] J. Kim, T. Noh, W. Oh, S. Park, An improved hybrid algorithm to bisection method and Newton-Raphson method, *Appl. Math. Sci.* 11 (2017), 2789–2797.
- [12] V. A. Patel, *Numerical Analysis*, Harcourt Brace College Publishers, Florida, 1994.
- [13] J. Naghipoor, S. A. Ahmadian, Ali R. Soheili, An improved regula falsi method for finding simple zeros of nonlinear equations, *Appl. Math. Sci.* 2 (2008), 381–386.
- [14] Y. Zhu, X. Wu, A free-derivative iteration method of order three having convergence of both point and interval for nonlinear equations, *Appl. Math. Comput.* 137 (2003), 49-55.