



Available online at <http://scik.org>

J. Semigroup Theory Appl. 2018, 2018:4

<https://doi.org/10.28919/jsta/3547>

ISSN: 2051-2937

## CONGRUENCES ON \*-BISIMPLE TYPE A I-SEMIGROUPS

NDUBUISI R.U.\* , ASIBONG-IBE U.I.

Department of Mathematics & Statistics, University of Port Harcourt, Port Harcourt, Nigeria

Copyright © 2018 Ndubuisi and Asibong-Ibe. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Abstract.** This paper studies congruences on a \*-bisimple type A I-semigroup in the light of known results in the areas of inverse semigroups and type A  $\omega$ -semigroups. It turns out that for a \*-bisimple type A I-semigroup, we have the idempotent-separating congruence and the minimum cancellative monoid congruence.

**Keywords:** type A I-semigroups; idempotent-separating; cancellative monoid congruence; generalized Bruck-Reilly \*-extension.

**2010 Mathematics Subject Classification:** 20M18.

### 1. Introduction and Summary

Let  $S$  be a semigroup and let  $E(S)$  denote the set of its idempotents. As well known,  $E(S)$  is partially ordered in the sense that: if  $e, f \in E(S)$ ,  $e \leq f$  if and only if  $ef = fe = e$ . Let  $I$  denote the set of all integers and let  $\mathbb{N}^0$  denote the set of nonnegative integers. A semigroup  $S$  is called an I-semigroup if and only if  $E(S)$  is order isomorphic to  $I$  under the reverse of the partial order. The \*-bisimple type A I-semigroup have been classified by Shang and Wang in [9]. The case in which  $\mathcal{D}^* = \tilde{\mathcal{D}}$  was shown to be the generalized Bruck-Reilly \*-extension of a cancellative monoid.

The main purpose of this paper is to present an explicit description of the congruences on \*-bisimple type A I-semigroups.

This work is divided into 5 sections; section 2 contains some preliminaries and results concerning \*-bisimple type A I-semigroups. The content of section 3 is the characterization of the idempotent-separating congruences on \*-bisimple type A I-semigroups. A description of the minimum cancellative monoid congruence on \*-bisimple type A I-semigroup is the subject of section 4 while the maximum idempotent-separating congruence is treated in section 5.

Now we recall some definitions which will be useful in the study. Terms not given here can be found in [4], [6] and [9], for more detailed knowledge.

---

\*Corresponding author

E-mail address: [u\\_ndubuisi@yahoo.com](mailto:u_ndubuisi@yahoo.com)

Received October 23, 2017

A semigroup  $S$  is said to be

- regular if all its elements are regular. Let  $S$  be a semigroup. An element  $x \in S$  is said to be regular if there exists  $y \in S$  such that  $xyx = x$ .
- unit regular if for each  $x \in S$  there exists a unit  $y$  of  $S$  for which  $x = xyx$ .
- An element  $x \in S$  is said to be coregular and  $y$  its coinverse if  $x = xyx = yxy$ .  $S$  is coregular if all its elements are coregular.
- orthodox if it is regular and the set  $E(S)$  of idempotent elements of the semigroup  $S$  forms a subsemigroup.

Let  $S$  be a semigroup and  $a, b \in S$ . The elements  $a$  and  $b$  in  $S$  are said to be  $\mathcal{R}^*$ -related written  $a \mathcal{R}^* b$  if and only if  $a$  and  $b$  are related in  $\mathcal{R}$  in some oversemigroup of  $S$ . Dually, we can define the relation  $\mathcal{L}^*$ . The following Lemma gives an alternative characterization of  $\mathcal{R}^*$ , the dual for the relation  $\mathcal{L}^*$ .

**Lemma 1.1** [4]. Let  $S$  be a semigroup and  $a, b \in S$ . Then  $a \mathcal{R}^* b$  if and only if for all  $x, y \in S^1$ ,  $xa = ya$  if and only if  $xb = yb$ .

As an easy but useful consequence of Lemma 1.1, we have

**Lemma 1.2** [4]. Let  $S$  be a semigroup and  $a, e^2 = e \in S$ . Then  $a \mathcal{R}^* e$  if and only if for any  $x, y \in S^1$ ,  $xa = ya$  implies  $xe = ye$ .

The join of the equivalence relations  $\mathcal{R}^*$  and  $\mathcal{L}^*$  is denoted by  $\mathcal{D}^*$  and their intersection by  $\mathcal{H}^*$ . Thus  $a \mathcal{H}^* b$  if and only if  $a \mathcal{R}^* b$  and  $a \mathcal{L}^* b$ . In general  $\mathcal{R}^* \circ \mathcal{L}^* \neq \mathcal{R}^* \circ \mathcal{L}^*$  (see [4]). Basically,  $a \mathcal{D}^* b$  if and only if there exist elements  $x_1, x_2, \dots, x_{2n-1}$  in  $S$  such that  $a \mathcal{L}^* x_1 \mathcal{R}^* x_2 \dots x_{2n-1} \mathcal{R}^* b$ .

Following Fountain [5] a semigroup is an abundant semigroup if every  $\mathcal{L}^*$ -class and every  $\mathcal{R}^*$ -class in  $S$  contain idempotents. An abundant semigroup  $S$  is an adequate [5] if  $E(S)$  forms a semilattice. In an adequate semigroup every  $\mathcal{L}^*$ -class  $\mathcal{R}^*$ -class contains a unique idempotent.

Let  $a$  be an element of an adequate semigroup  $S$ , and  $a^*$  ( $a^\dagger$ ) denotes the unique idempotent in the  $\mathcal{L}^*$ -class  $L_a^*$  ( $\mathcal{R}^*$ -class  $R_a^*$ ) containing  $a$ .

Fountain in [3] introduced the concept of right type A semigroup as special type of right PP monoids which is  $e$ -cancellable for an idempotent. He followed it in [4] with introduction of type a as an adequate semigroup satisfying certain internal conditions. An adequate semigroup  $S$  is a type A semigroup if  $ea = a(ea)^*$  and  $ae = (ae)^\dagger a$  for all  $a \in S$  and  $e \in E(S)$ . If a type A semigroup  $S$  contain precisely one  $\mathcal{D}^*$ -class it is said to be a  $*$ -bisimple type A semigroup.  $*$ -bisimple type A semigroup has been studied in [1].

## 2. The $*$ -Bisimple Type A I-Semigroup

In [9], Yu Shang and Limin Wang considered a similar construction of the one given earlier by Warne [10]. They used this construction to give the structure theorem for \*-bisimple type A I-semigroups. We now introduce the construction.

Let  $M$  be a monoid with  $\mathcal{H}_1^*$  as the  $\mathcal{H}^*$ -class which contain the identity element 1 of  $M$ . Let  $S = M \times I \times I$  (where  $I$  denotes the set of all integers) with multiplication defined by the rule

$$(x, m, n)(y, p, q) = \begin{cases} (x \cdot f_{n-p,p}^{-1} \cdot y \theta^{n-p} \cdot f_{n-p,q}, m, n+q-p) & \text{if } n \geq p \\ (f_{p-n,n}^{-1} \cdot x \theta^{p-n} \cdot f_{p-n,n} \cdot y, m+p-n, q) & \text{if } n \leq p \end{cases}$$

where  $\theta$  is an endomorphism of  $M$  with images in  $\mathcal{H}_1^*$ .  $\theta^0$  denotes the identity automorphism of  $M$ , and for  $m \in \mathbb{N}^0, n \in I, f_{0,n} = 1$ , the identity of  $M$ , and for  $m > 0, f_{m,n} = u_{n+1} \theta^{m-1} \cdot u_{n+2} \theta^{m-2} \dots u_{n+(m-1)} \theta \cdot u_{n+m}$  and  $f_{m,n}^{-1} = u_{n+m}^{-1} \cdot u_{n+(m-1)}^{-1} \theta \dots u_{n+2}^{-1} \theta^{m-2} \cdot u_{n+1}^{-1} \theta^{m-1}$ , where  $\{u_n \mid n \in I\}$  is a collection of elements of  $H_1$  with  $u_n = 1$  if  $n > 0$ .

Under the above multiplication,  $S = M \times I \times I$  is a semigroup (see [9]) and this semigroup is called the generalized Bruck-Reilly \*-extensions of  $M$  determined by  $\theta$  and it is usually denoted by  $S = GBR^*(M, \theta)$ .

The following results are proved in [9]. We give a sketch proof of (i), (iii) and (v)

**Lemma 2.1.** Let  $(x, m, n), (y, p, q) \in GBR^*(M, \theta)$ . Then

- (i)  $(x, m, n) \mathcal{L}^* (y, p, q)$  if and only if  $n = q$  and  $x \mathcal{L}^*(M) y$ .
- (ii)  $(x, m, n) \mathcal{R}^* (y, p, q)$  if and only if  $m = p$  and  $x \mathcal{R}^*(M) y$ .
- (iii)  $(x, m, n) \in E(GBR^*(M, \theta))$  if and only if  $m = n$  and  $x \in E(M)$ .
- (iv)  $(x, m, n)$  has an inverse  $(y, p, q) \in S$  if and only if  $p = n, q = m$  and  $x$  is the inverse of  $y \in M$ .
- (v)  $GBR^*(M, \theta)$  is a type A semigroup if and only if  $M$  is a type A semigroup.

**Proof.** (i) Let  $(x, m, n) \mathcal{L}^* (y, p, q)$ . For  $(e, 0, 0), (e, n, n) \in GBR^*(M, \theta)$  we have

$$(x, m, n)(e, 0, 0) = (x, m, n)(e, n, n),$$

$$(y, p, q)(e, 0, 0) = (y, p, q)(e, n, n).$$

Consequently,

$$(y, p, q) = (y, p, q)(e, n, n). \text{ If } q < n, \text{ this gives}$$

$$(y, p, q) = (f_{n-q,p}^{-1} \cdot y \theta^{n-q} \cdot f_{n-q,q} \cdot e, p+n-q, n).$$

Comparing the third coordinates gives  $q = n$ , which is a contradiction. Thus  $q \geq n$ .

Similarly, using the idempotent  $(e, q, q)$  we have

$$(x, m, n)(e, q, q) = \begin{cases} (x \cdot f_{n-q,q}^{-1} \cdot e\theta^{n-q} \cdot f_{n-q,q}, m, n+q-q) & \text{if } n \geq q \\ (f_{q-n,m}^{-1} \cdot x\theta^{q-n} \cdot f_{q-n,n} \cdot e, m+q-n, q) & \text{if } n \leq q \end{cases}$$

So we deduce that  $q \leq n$  and so  $q = n$ .

Conversely, let  $n = q$ . For any arbitrary elements  $(v, i, j), (w, l, k) \in GBR^*(M, \theta)$ ,

$$(x, m, n)(v, i, j) = (x, m, n)(w, l, k).$$

Suppose  $n \geq i$  and  $n \geq l$ . Then

$$(x \cdot f_{n-i,i}^{-1} \cdot v\theta^{n-i} \cdot f_{n-i,j}, m, n+j-i) = (x \cdot f_{n-l,l}^{-1} \cdot w\theta^{n-l} \cdot f_{n-l,k}, m, n+k-l).$$

Comparing the first and the third coordinates gives

$$x \cdot f_{n-i,i}^{-1} \cdot v\theta^{n-i} \cdot f_{n-i,j} = x \cdot f_{n-l,l}^{-1} \cdot w\theta^{n-l} \cdot f_{n-l,k} \quad \text{and} \quad n+j-i = n+k-l.$$

This implies

$$y \cdot f_{n-i,i}^{-1} \cdot v\theta^{n-i} \cdot f_{n-i,j} = y \cdot f_{n-l,l}^{-1} \cdot w\theta^{n-l} \cdot f_{n-l,k} \quad \text{and} \quad n+j-i = n+k-l.$$

Hence,  $(y, p, n)(v, i, j) = (y, p, n)(w, l, k)$ .

(ii) The proof is similar to the proof of (i).

(iii) Let  $(x, m, n) \in E(GBR^*(M, \theta))$ . Then

$$\begin{aligned} (x, m, n) &= (x, m, n)(x, m, n) && = \\ &\begin{cases} (x \cdot f_{n-m,m}^{-1} \cdot x\theta^{n-m} \cdot f_{n-m,n}, m, n+n-m) & \text{if } n \geq m \\ (f_{m-n,m}^{-1} \cdot x\theta^{m-n} \cdot f_{m-n,n} \cdot x, m+m-n, n) & \text{if } n \leq m \end{cases} \end{aligned}$$

thus  $m = n$  and  $x^2 = x$ .

Conversely, let  $m = n$  and  $x \in E(M)$ . Then certainly  $(x, m, n)(x, m, n) = (x, m, n)$ . From which it follows that  $(x, m, n) \in E(GBR^*(M, \theta))$ .

(iv) The proof is clear.

(v) We only prove that  $GBR^*(M, \theta)$  is right type A, as the proof that  $GBR^*(M, \theta)$  is left type A is dual.

Let  $(e, m, m), (e, n, n) \in E(GBR^*(M, \theta))$ . Suppose that  $m > n$ . Then

$$\begin{aligned} (e, m, m), (e, n, n) &= (e \cdot f_{m-n,n}^{-1} \cdot e\theta^{m-n} \cdot f_{m-n,n}, m) \\ &= (f_{m-n,n}^{-1} \cdot e\theta^{m-n} \cdot f_{m-n,n} \cdot e, m, m) \\ &= (e, n, n)(e, m, m). \end{aligned}$$

Thus the idempotents of  $GBR^*(M, \theta)$  commute. So every  $\mathcal{L}^*$ -class of  $GBR^*(M, \theta)$  contain an idempotent.

Let  $(x, p, q) \in GBR^*(M, \theta)$ . Suppose  $m \geq p$ . Then

$$\begin{aligned} (x, p, q)[(e, m, m)(x, p, q)]^* &= (x, p, q)(e \cdot f_{m-p,p}^{-1} \cdot x\theta^{m-p} \cdot f_{m-p,q}, m, m+q-p)^* \\ &= (x, p, q)(e \cdot f_{m-p,p}^{-1} \cdot x\theta^{m-p} f_{m-p,q}, m+q-p, m+q-p) \\ &= (e, m, m)(x, p, q). \end{aligned}$$

**Theorem 2.2** (Structure theorem)

Let  $S = GBR^*(M, \theta)$  be the generalized Bruck-Reilly \*-extensions of  $M$  determined by  $\theta$ . Then  $S$  is a \*-bisimple type A I-semigroup. Conversely, every \*-bisimple type A I-semigroup is isomorphic to  $GBR^*(M, \theta)$ .

**Proof.** It is known that  $S = GBR^*(M, \theta)$  is a type A semigroup. That  $S$  is \*-bisimple follows from Lemma 2.1 (i) & (ii).

Next, let  $e_m = (e, m, m)$  and  $e_n = (e, n, n) \in E(S)$ . Then for  $m \geq n$ .

$$\begin{aligned} e_m e_n &= (e, m, m)(e, n, n) = (e \cdot f_{m-n,n}^{-1} \cdot e\theta^{m-n} \cdot f_{m-n,n}, m, m+n-n) \\ &= (e, m, m) = e_m \\ &= (e, n, n)(e, m, m) = e_n e_m \end{aligned}$$

Thus  $e_m \leq e_n$  if and only if  $m \geq n$ , which shows that  $E(S)$  is a chain

$$\dots > (e, -2, -2) > (e, -1, -1) > (e, 0, 0) > (e, 1, 1) > (e, 2, 2) > \dots$$

Hence  $S$  is a \*-bisimple type A I-semigroup. The converse of the proof is a routine check.

From Lemma 2.1(iv), we have the following result

**Corollary 2.3.** Let  $M$  be a monoid. Then  $S = GBR^*(M, \theta)$  is regular if and only if  $M$  is regular.

The following results show some other properties of  $S = GBR^*(M, \theta)$ .

**Proposition 2.4.** Let  $S = GBR^*(M, \theta)$ . Then  $S$  is unit regular if and only if  $M$  is unit regular.

**Proof.** Let  $S = GBR^*(M, \theta)$  be unit regular. Then for any  $(x, m, n) \in S$ , there exists an element  $(y, n, m) \in G$  (where  $G$  is the group of units of  $GBR^*(M, \theta)$ ) such that

$$(x, m, n)(y, n, m)(x, m, n) = (x, m, n).$$

By considering left-hand side of the equation, we get

$$\begin{aligned} (x, m, n)(y, n, m)(x, m, n) &= ((x, m, n)(y, n, m))(x, m, n) \\ &= (x \cdot f_{n-n,n}^{-1} \cdot y\theta^{n-n} \cdot f_{n-n,m}, m, n+m-n)(x, m, n) \\ &= (xy, m, m)(x, m, n) = (xyx, m, n). \end{aligned}$$

Therefore we obtain  $x = xyx$ . Consequently,  $M$  is unit regular.

Conversely, let us suppose that  $M$  is unit regular. Then for  $x \in M$ , there exists an element  $x \in G_M$  (where  $G_M$  is the group of units of  $M$ ) such that obtain  $x = xyx$ . Now we need to show that for any  $(x, m, n) \in GBR^*(M, \theta)$ , there exist an element  $(y, p, q) \in G_M$  such that

$$(x, m, n) = (x, m, n)(y, p, q)(x, m, n).$$

Here we take  $p = n$ ,  $q = m$ , then we have  $(x, m, n)(y, n, m)(x, m, n) = (xyx, m, n)$ . Since we have  $x = xyx$ , for any  $x \in M, y \in G_M$ , we obtain  $(x, m, n)(y, p, q)(x, m, n) = (x, m, n)$ . Thus  $S$  is unit regular.

**Proposition 2.5** Let  $M$  be a monoid. Then  $M' = \{(x, m, m) \mid x \in M, m \in \mathbb{N}^0\} \leq GBR^*(M, \theta)$  is coregular if and only if  $M$  is coregular.

**Proof.** Let  $M' \leq GBR^*(M, \theta)$  be coregular. Then for  $(x, 0, 0) \in GBR^*(M, \theta)$ , there exists an element  $(y, n, n) \in GBR^*(M, \theta)$  such that

$$((x, 0, 0)(y, n, n))(x, 0, 0) = (xyx, n, n) = (x, 0, 0) \quad (1)$$

$$((y, n, n)(x, 0, 0))(y, n, n) = (yxy, n, n) = (x, 0, 0) \quad (2)$$

From (1) and (2), we have that  $n = 0$ ,  $xyx = x$  and  $yxy = x$ . Thus  $M$  is coregular.

Conversely, let  $M$  be coregular. Then there exists  $y \in M$ , with  $xyx = x$  and  $yxy = x$ . Thus for  $(x, m, n) \in GBR^*(M, \theta)$ , we have

$$\begin{aligned} ((x, m, n)(y, m, m))(x, m, m) &= (xy, m, m)(x, m, m) \\ &= (xyx, m, m) \\ &= (x, m, m). \end{aligned}$$

$$\begin{aligned} ((y, m, m)(x, m, m))(y, m, m) &= (yx, m, m)(y, m, m) \\ &= (yxy, m, m) \\ &= (x, m, m). \end{aligned}$$

Therefore,  $M' = \{(x, m, m) \mid x \in M, m \in \mathbb{N}^0\} \leq GBR^*(M, \theta)$  is coregular.

It is important to note that not all regular semigroups are coregular. This is shown in the example below.

**Example 2.6.** Let  $X$  and  $Y$  be non-empty sets and set  $T = X \times Y$  with the binary operation

$$(x, y)(u, v) = (x, v), \text{ for all } x, u \in X, y, v \in Y.$$

It can be easily seen that  $T$  is a semigroup. This semigroup is called a rectangular band.  $T$  is also regular, since for  $(x, y), (u, v) \in T$  we have  $(x, y)(u, v)(x, y) = (x, y)$ .

To show that  $T$  is not coregular, let  $(x, y), (u, v) \in T$ , then we have

$$(x, y)(u, v)(x, y) = (x, y),$$

$$(u, v)(x, y)(u, v) = (u, v).$$

So  $(x, y) \neq (u, v)$ . Thus  $T$  is not coregular.

In the next theorem, we consider the orthodox property of  $GBR^*(M, \theta)$

**Theorem 2.7.** Let  $S = GBR^*(M, \theta)$ . Then  $S$  is orthodox if and only if  $M$  is orthodox.

**Proof.** Let  $GBR^*(M, \theta)$  be orthodox. By Corollary 2.3, we know that  $M$  is regular. Then it remains to show that  $E(M)$  is a subsemigroup of  $M$ . In particular for each  $e, e' \in E(M)$ ,

$$\begin{aligned} (e, m, m)(e', m, m) &= (e \cdot f_{m-m, m}^{-1} \cdot e' \theta^{m-m} \cdot f_{m-m, m}, m, m + m - m) \\ &= (ee', m, m) \end{aligned}$$

is an idempotent of  $GBR^*(M, \theta)$  and so  $(ee')^2 = ee'$ . Hence  $M$  is orthodox.

Conversely, let  $M$  be orthodox. Then  $M$  is regular, and  $E(M)$  is a subsemigroup of  $M$ . By Corollary 2.3, we know that  $GBR^*(M, \theta)$  is regular.

Next, we show that  $(e, m, m)(e', n, n) \in E(GBR^*(M, \theta))$ . From the multiplication  $(e, m, m)(e', n, n)$ , we have the following cases:

Case (1): If  $m \geq n$ , we have

$$\begin{aligned} (e, m, m)(e', n, n) &= \left( (e \cdot f_{m-n, n}^{-1}) \cdot (e' \theta^{m-n} \cdot f_{m-n, n}), m, m + n - n \right) \\ &= \left( (e \cdot f_{m-n, n}^{-1}) \cdot (e' \theta^{m-n} \cdot f_{m-n, n}), m, m \right). \end{aligned}$$

Since  $e, e' \in E(M)$ , we deduce that  $e \cdot f_{m-n, n}^{-1}, e' \theta^{m-n} \cdot f_{m-n, n} \in E(M)$ . But the idempotents in  $M$  are commutative, consequently

$$(e \cdot f_{m-n, n}^{-1}) \cdot (e' \theta^{m-n} \cdot f_{m-n, n}) = (e' \theta^{m-n} \cdot f_{m-n, n}) \cdot (e \cdot f_{m-n, n}^{-1}).$$

So  $(e' \theta^{m-n} \cdot f_{m-n, n}), (e \cdot f_{m-n, n}^{-1}) \in E(GBR^*(M, \theta))$ . Therefore  $E(GBR^*(M, \theta))$  is a subsemigroup of  $GBR^*(M, \theta)$ .

Case (2): If  $m \leq n$ , we have

$$\begin{aligned} (e, m, m)(e', n, n) &= \left( (f_{n-m, n}^{-1} \cdot e \theta^{n-m}) \cdot (f_{n-m, m} \cdot e'), m + n - m, n \right) \\ &= \left( (f_{n-m, m}^{-1} \cdot e \theta^{n-m}) \cdot (f_{n-m, m} \cdot e'), n, n \right). \end{aligned}$$

From here, since  $(f_{n-m, m}^{-1} \cdot e \theta^{n-m}), (f_{n-m, m} \cdot e') \in E(M)$  and the idempotents in  $M$  are commutative, we deduce that  $E(GBR^*(M, \theta))$  is a subsemigroup of  $GBR^*(M, \theta)$ .

The connection between the Green's \*-relations and congruences lies on the fact that  $\mathcal{L}^*$  is a right congruence and  $\mathcal{R}^*$  is a left congruence. It can be easily verified that  $\mathcal{H}^*$  is a congruence on  $S = GBR^*(M, \theta)$ . In our next section, we shall characterize the congruences on  $S = GBR^*(M, \theta)$ .

### 3. Idempotent-separating congruences

The following terms adopted from [8] will be used in the description of congruences on \*-bisimple type A I-semigroups.

**Definition 3.1.** Let  $S = GBR^*(M, \theta)$  be a \*-bisimple type A I-semigroup where  $\theta : M \rightarrow \mathcal{H}_1^*$ . Let  $\mathcal{H}^* = \rho$  be a congruence on  $S$ . Let us use  $\rho(M)$  to denote the congruence on  $M$  induced by  $\rho$ , via the restriction of  $\rho$  to the monoid  $\{(x, 0, 0) : x \in M\}$ .

**Definition 3.2.** A congruence  $\gamma$  on  $M$  is said to be  $\theta$ -admissible if  $x \gamma y$  implies  $x\theta \gamma y\theta$ , for any  $x, y \in M$ .

A typical idempotent-separating congruence on  $S = GBR^*(M, \theta)$  is characterized as follows:

**Theorem 3.3.** Let  $S = GBR^*(M, \theta)$  be a  $*$ -bisimple type A I-semigroup and let  $\rho$  be a congruence on  $S = GBR^*(M, \theta)$ . Then  $\rho(M)$  is  $\theta$ -admissible. Conversely, if  $\gamma$  is any  $\theta$ -admissible congruence on  $M$ , then the relation on  $S$  defined by

$$[(x, m, n)(y, p, q)] \in \gamma(S) \text{ if and only if } m = p, n = q \text{ and } (x, y) \in \gamma$$

is an idempotent-separating congruence.

**Proof.** Suppose  $x \rho(M) y$ . Then we have that  $(x, 0, 0) \rho (y, 0, 0)$ .

Consequently,

$$(x, 0, 0)(e, 1, 1) \rho (y, 0, 0)(e, 1, 1).$$

But  $(x, 0, 0)(e, 1, 1) = (f_{1,0}^{-1} \cdot x \theta f_{1,0} \cdot e, 1, 1)$  and  $(y, 0, 0)(e, 1, 1) = (f_{1,0}^{-1} \cdot y \theta f_{1,0} \cdot e, 1, 1)$ .

Thus  $(f_{1,0}^{-1} \cdot x \theta f_{1,0} \cdot e, 1, 1) \rho (f_{1,0}^{-1} \cdot y \theta f_{1,0} \cdot e, 1, 1) = (x \theta, 1, 1) \rho (y \theta, 1, 1)$ .

Since  $(x \theta, 1, 1) \rho (y \theta, 1, 1)$ , then  $(x \theta, 1, 1) = (y \theta, 1, 1)$ .

Also we have  $(e, 0, 1)(x \theta, 1, 1)(e, 1, 0) \rho (e, 0, 1)(y \theta, 1, 1)(e, 1, 0)$ .

But  $(e, 0, 1)(x \theta, 1, 1)(e, 1, 0) = (x \theta, 0, 0)$  and  $(e, 0, 1)(y \theta, 1, 1)(e, 1, 0) = (y \theta, 0, 0)$ .

Thus  $(x \theta, 0, 0) \rho (y \theta, 0, 0)$ . Since  $(x \theta, 0, 0) \rho (y \theta, 0, 0)$ , then  $x \theta \rho(M) y \theta$ .

Conversely, let  $\gamma$  be a  $\theta$ -admissible congruence on  $M$ . We first show that  $\gamma(S)$  is an equivalence relation.

$[(x, m, n)(x, m, n)] \in \gamma(S)$  since  $(x, x) \in \gamma$ . Thus  $\gamma(S)$  is reflexive. By definition,  $\gamma(S)$  is symmetric.

To show transitivity, let  $(x, m, n) \gamma(S) (y, p, q)$  and  $(y, p, q) \gamma(S) (z, i, j)$  for all  $(x, m, n), (y, p, q), (z, i, j) \in S$ . Then we have  $m = p, n = q, (x, y) \in \gamma$  and  $p = i, q = j, (y, z) \in \gamma$ .

Consequently,  $m = i, n = j$ . Hence  $(x, z) \in \gamma$ , which means that  $\gamma(S)$  is transitive.

Next is to show that  $\gamma(S)$  is a congruence. Now let  $a = (x, m, n), b = (y, p, q)$ . That  $\gamma(S)$  is a congruence entails showing that

$$a \gamma(S) b \text{ implies } ax \gamma(S) bx \quad (\text{for right congruence})$$

$$a \gamma(S) b \text{ implies } xa \gamma(S) xb \quad (\text{for left congruence})$$

$\forall x = (z, k, l) \in S = GBR^*(M, \theta)$ .

Consequently,

$$ax = (x, m, n)(z, k, l) = \begin{cases} (x \cdot f_{n-k,k}^{-1} \cdot z \theta^{n-k} \cdot f_{n-k,l}, m, n + l - k) & \text{if } n \geq k \\ (f_{k-n,m}^{-1} \cdot x \theta^{k-n} \cdot f_{k-n,n} \cdot z, m + k - n, l) & \text{if } n \leq k \end{cases}$$

$$bx = (y, p, q)(z, k, l) = \begin{cases} (y \cdot f_{q-k,k}^{-1} \cdot z \theta^{q-k} \cdot f_{q-k,l}, p, q + l - k) & \text{if } q \geq k \\ (f_{k-q,p}^{-1} \cdot y \theta^{k-q} \cdot f_{k-q,q} \cdot z, p + k - q, l) & \text{if } q \leq k \end{cases}$$

So if  $(x, m, n) \gamma(S) (y, p, q)$ , then

$$(x, m, n)(z, k, l) \gamma(S) (y, p, q)(z, k, l) =$$



## CONGRUENCES ON \*-BISIMPLE TYPE A I-SEMIGROUPS

$$\gamma(S) \begin{cases} (x \cdot f_{n-k,k}^{-1} \cdot z\theta^{n-k} \cdot f_{n-k,l}, m, n+l-k) & \text{if } n \geq k \\ (f_{k-n,m}^{-1} \cdot x\theta^{k-n} \cdot f_{k-n,n} \cdot z, m+k-n, l) & \text{if } n \leq k \end{cases}$$

$$\gamma(S) \begin{cases} (y \cdot f_{q-k,k}^{-1} \cdot z\theta^{q-k} \cdot f_{q-k,l}, p, q+l-k) & \text{if } q \geq k \\ (f_{k-q,p}^{-1} \cdot y\theta^{k-q} \cdot f_{k-q,q} \cdot z, p+k-q, l) & \text{if } q \leq k \end{cases}$$

But  $(x, m, n) \gamma(S) (y, p, q)$  if and only if  $m = p, n = q$  and  $x \gamma y$ .

Thus, we have that

$$\begin{cases} (x \cdot f_{n-k,k}^{-1} \cdot z\theta^{n-k} \cdot f_{n-k,l}, m, n+l-k) & \text{if } n \geq k \\ (f_{k-n,m}^{-1} \cdot x\theta^{k-n} \cdot f_{k-n,n} \cdot z, m+k-n, l) & \text{if } n \leq k \end{cases}$$

$$\gamma(S) \begin{cases} (y \cdot f_{n-k,k}^{-1} \cdot z\theta^{n-k} \cdot f_{n-k,l}, m, n+l-k) & \text{if } n \geq k \\ (f_{k-n,m}^{-1} \cdot y\theta^{k-n} \cdot f_{k-n,n} \cdot z, m+k-n, l) & \text{if } n \leq k \end{cases}$$

Hence  $\gamma(S)$  is a right congruence.

That  $\gamma(S)$  is a left congruence follows similarly. Thus  $\gamma(S)$  is a congruence.

Futhermore,  $(e, m, m) \gamma(S) (e, n, n) \Rightarrow m = n$  which implies that  $(e, m, m) = (e, n, n)$ . Thus  $\gamma(S)$  is an idempotent-separating congruence.

**Remark 3.4.**  $\mathcal{H}^*$  is an idempotent-separating congruence on  $S = GBR^*(M, \theta)$  and  $\gamma(S) \subseteq \mathcal{H}^*$ .

#### 4. Minimum cancellative monoid congruence

The idea of the minimum cancellative monoid congruence is to obtain a congruence  $\sigma$  on  $S$ , a type A semigroup with respect to which  $S/\sigma$  is cancellative.

Here we will determine the minimum cancellative monoid congruence on  $S = GBR^*(M, \theta)$ , as follows:

Now let  $(h, m, n), (x, i, j) \in S = GBR^*(M, \theta)$ . Define a relation  $\sigma$  on  $S = GBR^*(M, \theta)$  by the rule

$$(h, m, n) \sigma (x, i, j) \text{ if and only if } m - n = i - j, h\theta^i = x\theta^m \text{ and } x\theta^i = h\theta^m.$$

**Lemma 4.1.**  $\sigma$  is a congruence on  $S$ .

**Proof.** That  $\sigma$  is symmetric and reflexive is known. To show that  $\sigma$  is transitive, let  $(h, m, n) \sigma (x, i, j)$  and  $(x, i, j) \sigma (y, p, q)$  for  $(h, m, n), (x, i, j), (y, p, q) \in S$ . Then  $m - n = i - j$  and  $i - j = p - q$  and so  $m - n = p - q$ .

Consequently,  $x\theta^i = h\theta^m$  and  $y\theta^p = x\theta^i$  implies  $y\theta^p = h\theta^m$ .

Also  $h\theta^i = x\theta^m$  and  $x\theta^p = y\theta^i$  implies that  $h\theta^i = (y\theta^{i-p})\theta^m = y\theta^{i-p+m}$ . Then  $h\theta^{i+p} = y\theta^{i-p+m+p} = y\theta^{i+m}$ . Hence  $h\theta^p = y\theta^m$  which shows that  $\sigma$  is transitive.

Next we show that  $\sigma$  is a congruence. Now let  $u = (h, m, n)$ ,  $v = (x, i, j)$ . That  $\sigma$  is a congruence we show that  $\sigma$  is both a left and right congruence. That is

$$\forall z \in S, \quad u \sigma v \implies uz \sigma vz \quad (\text{for right congruence})$$

and

$$\forall z \in S \quad u \sigma v \implies zu \sigma zv \quad (\text{for left congruence}).$$

Let  $z = (y, p, q) \in S$ . Then

$$uz = (h, m, n)(y, p, q) = \begin{cases} (h \cdot f_{n-p,p}^{-1} \cdot y\theta^{n-p} \cdot f_{n-p,q}, m, n+q-p) & \text{if } n \geq p \\ (f_{p-n,n}^{-1} \cdot h\theta^{p-n} \cdot f_{p-n,n} \cdot y, m+p-n, q) & \text{if } n \leq p \end{cases}$$

and

$$vz = (x, i, j)(y, p, q) = \begin{cases} (x \cdot f_{j-p,p}^{-1} \cdot y\theta^{j-p} \cdot f_{j-p,q}, i, j+q-p) & \text{if } j \geq p \\ (f_{p-j,j}^{-1} \cdot x\theta^{p-j} \cdot f_{p-j,j} \cdot y, i+p-j, q) & \text{if } j \leq p \end{cases}$$

Evidently if  $(h, m, n) \sigma (x, i, j)$ , we have

$$m - (n + q - p) = (m - n) + (p - q) \quad \text{and} \quad i - (j + q - p) = (i - j) + (p - q)$$

$$m + p - n - q = (m - n) + (p - q) \quad \text{and} \quad i + p - j - q = (i - j) + (p - q).$$

But  $m - n = i - j$  and so  $(m - n) + (p - q) = (i - j) + (p - q)$ .

For the first outer part, we know from definition that  $h\theta^i = x\theta^m$  and  $h\theta^n = x\theta^j$ . It suffices to show that  $(h\theta^{p-n} \cdot y)\theta^{i+p-j} = (x\theta^{p-j} \cdot y)\theta^{m+p-n}$ .

Considering the left hand side of the equation we have

$$\begin{aligned} (h\theta^{p-n} \cdot y)\theta^{i+p-j} &= h\theta^{p+p+i-n-j} \cdot y\theta^{p-j+i} \\ &= h\theta^{i+(p+p)-j-n} \cdot y\theta^{i+p-j} \\ &= (h\theta^i)\theta^{p-j-n+p} \cdot y\theta^{p+(i-j)} \end{aligned}$$

But  $i - j = m - n$  and  $h\theta^i = x\theta^m$ .

$$\begin{aligned} \text{Therefore, } (h\theta^i)\theta^{p-j-n+p} \cdot y\theta^{p+(i-j)} &= (x\theta^m)\theta^{p-j-n+p} \cdot y\theta^{p+(m-n)} \\ &= x\theta^{m+p+p-j-n} \cdot y\theta^{p+m-n} \\ &= (x\theta^{p-j} \cdot y)\theta^{m+p-n} \end{aligned}$$

as required.

Hence  $\sigma$  is a right congruence. That  $\sigma$  is a left congruence follows similarly. Consequently  $\sigma$  is a congruence.

**Lemma 4.2.**  $\sigma$  is a cancellative monoid.

**Proof.** Since  $(e, m, m) \sigma (e, n, n)$  for  $m, n \in I$ , it follows that the class of  $\sigma$  containing the idempotents is the identity element for  $S/\sigma$ . Thus  $(1, m, n) \sigma (y, p, q) \sigma = (y, p, q) \sigma$  and hence  $S/\sigma$  is a monoid.

Next is to show that  $S/\sigma$  is cancellative. Now let  $u = (h, m, n), v = (x, i, j)$ .

That  $S/\sigma$  is cancellative entails showing that for all  $z \in S$ ,

$$u \sigma z \sigma = v \sigma z \sigma \Rightarrow u \sigma = v \sigma \quad (\text{for right cancellative})$$

and

$$z \sigma u \sigma = z \sigma v \sigma \Rightarrow u \sigma = v \sigma \quad (\text{for left cancellative}).$$

Let  $z = (y, p, q) \in S$ . Then

$$\begin{aligned} u \sigma z \sigma &= (h, m, n) \sigma (y, p, q) \sigma = (x, i, j) \sigma (y, p, q) \sigma \\ &= v \sigma z \sigma . \end{aligned}$$

Consequently,

$$\begin{aligned} &(h, m, n) \sigma (y, p, q) \sigma = (x, i, j) \sigma (y, p, q) \sigma \\ \Leftrightarrow &[(h, m, n)(y, p, q)] \sigma = [(x, i, j)(y, p, q)] \sigma \\ \Leftrightarrow &\begin{cases} (h \cdot f_{n-p,p}^{-1} \cdot y \theta^{n-p} \cdot f_{n-p,q} \cdot m, n + q - p) & \text{if } n \geq p \\ (f_{p-n,n}^{-1} \cdot h \theta^{p-n} \cdot f_{p-n,n} \cdot y, m + p - n, q) & \text{if } n \leq p \end{cases} \times \sigma \\ = &\begin{cases} (x \cdot f_{j-p,p}^{-1} \cdot y \theta^{j-p} \cdot f_{j-p,q} \cdot i, j + q - p) & \text{if } j \geq p \\ (f_{p-j,i}^{-1} \cdot x \theta^{p-j} \cdot f_{p-j,j} \cdot y, i + p - j, q) & \text{if } j \leq p \end{cases} \times \sigma \\ \Leftrightarrow &m - (n + q - p) = i - (y + q - p), (m + p - n) - q = (i + p - j) - q \end{aligned}$$

and

$$\begin{aligned} &(h \theta^{p-n} \cdot y) \theta^{i+p-j} = (x \theta^{p-j} \cdot y) \theta^{m+p-n} \\ \Leftrightarrow &(m - n) + (p - q) = (i - j) + (p - q) \end{aligned}$$

and

$$\begin{aligned} &h \theta^{p-n+(i-j)+p} \cdot y \theta^{p+(i-j)} = x \theta^{p-j+(m-n)+p} \cdot y \theta^{p+(m-n)} \\ \Leftrightarrow &m - n = i - j \text{ and } (h \theta^i)^{p+p-n-j} = (x \theta^m) \theta^{p+p-n-j} \\ \Leftrightarrow &m - n = i - j \text{ and } h \theta^i = x \theta^m \\ \Leftrightarrow &(h, m, n) \sigma (x, i, j) \end{aligned}$$

which shows that  $S/\sigma$  is right cancellative. That  $S/\sigma$  is left cancellative follows similarly, and we conclude that  $S/\sigma$  is cancellative.

**Lemma 4.3**  $\sigma$  is a minimum congruence.

**Proof.** Let  $\Gamma$  be any other cancellative monoid congruence. Then  $(1, n, n) \Gamma (1, 0, 0)$  for all  $n \in I$ . Suppose  $(h, m, n) \sigma (x, i, j)$ . Then we have from  $(h, m, n)(1, p, p) = (x, i, j)(1, p, p)$  for some  $p \in I$ ,

$$\begin{aligned}
& \Rightarrow \begin{cases} (h \cdot f_{n-p,p}^{-1} \cdot 1\theta^{n-p} \cdot f_{n-p,p}, m, n+p-p) & \text{if } n \geq p \\ (f_{p-n,m}^{-1} \cdot h\theta^{p-n} \cdot f_{p-n,n} \cdot 1, m+p-n, p) & \text{if } n \leq p \end{cases} \\
& = \begin{cases} (x \cdot f_{j-p,p}^{-1} \cdot 1\theta^{j-p} \cdot f_{j-p,p}, i, j+p-p) & \text{if } j \geq p \\ (f_{p-j,i}^{-1} \cdot x\theta^{p-j} \cdot f_{p-j,j} \cdot 1, i+p-j, p) & \text{if } j \leq p \end{cases} \\
& \Rightarrow \begin{cases} (h, m, n) & \text{if } n \geq p \\ (h\theta^{p-n}, m+p-n, p) & \text{if } n \leq p \end{cases} = \begin{cases} (x, i, j) & \text{if } j \geq p \\ (x\theta^{p-j}, i+p-j, p) & \text{if } j \leq p \end{cases}
\end{aligned}$$

But  $(1, n, n) \Gamma (1, 0, 0)$ , so  $(h, m, n)(1, p, p) \Gamma (h, m, n)$ .

Also,  $(x, i, j)(1, p, p) \Gamma (x, i, j)$ . Therefore  $(h, m, n) \Gamma (x, i, j)$ . Thus  $\sigma \subseteq \Gamma$ .

Combining Lemma 4.1 to Lemma 4.3, we have proved the following theorem:

**Theorem 4.4.** Let  $S = GBR^*(M, \theta)$  be a \*-bisimple type A I-semigroup and let  $\sigma$  be defined on  $S$  by the rule that  $(h, m, n) \sigma (x, i, j)$  if and only if  $m - n = i - j$ ,  $h\theta^i = x\theta^m$  and  $x\theta^i = h\theta^m$ . Then  $\sigma$  is the minimum cancellative monoid congruence on  $S$ .

## 5. The congruence $\mu$

Here we will determine the maximum congruence  $\mu$  on  $S = GBR^*(M, \theta)$  contained in  $\mathcal{H}^*$  by utilizing the approach of El-Qallali and Fountain [2].

Now let  $(e, m, m)$  and  $(e, n, n)$  be the idempotents in the  $\mathcal{R}^*$ -class and  $\mathcal{L}^*$ -class respectively. We define the relations  $\mu_R$  and  $\mu_L$  on  $S = GBR^*(M, \theta)$  as follows:

$$(x, m, n) \mu_L (y, p, q) \text{ if and only if } (e, n, n)(x, m, n) \mathcal{L}^* (e, n, n)(y, p, q), m - n = p - q, \\ x\theta^{n-m} = y\theta^{n-p} \text{ and } e\theta^{m-n} \cdot x = e\theta^{p-n} \cdot y.$$

$$(x, m, n) \mu_R (y, p, q) \text{ if and only if } (x, m, n)(e, m, m) \mathcal{R}^* (y, p, q)(e, m, m), m - n = p - q, \\ x\theta^{m-n} = y\theta^{m-q} \text{ and } x \cdot e\theta^{n-m} = y \cdot e\theta^{q-m}.$$

Consequently,

$$\mu = \mu_L \cap \mu_R.$$

With the above relation, we obtain the following results

**Proposition 5.1.** Let  $S = GBR^*(M, \theta)$ . Then  $\mu_L$  is the maximum congruence on  $S$  contained in  $\mathcal{L}^*$ .

**Proof.** Obviously,  $\mu_L$  is an equivalence on  $S$ . Since  $\mathcal{L}^*$  is a right congruence on  $S$ ,  $\mu_L$  is right compatible under the semigroup multiplication.

Next is to show that  $\mu_L$  is also left compatible under the semigroup multiplication. Now let  $(x, m, n), (y, p, q), (e, 0, 0) \in S$ . That  $\mu_L$  is left compatible entails showing that

$$(e, n, n)(x, m, n) \mathcal{L}^* (e, n, n)(y, p, q) \text{ implies } (e, 0, 0)(e, n, n)(x, m, n) \mathcal{L}^* (e, 0, 0)(e, n, n)(y, p, q).$$

Thus we have

$$(e, 0, 0)(e, n, n)(x, m, n) = (e\theta^n, n, n)(x, m, n),$$

and

$$(e, 0, 0)(e, n, n)(y, p, q) = (e\theta^n, n, n)(y, p, q).$$

Consequently,

$$(e\theta^n, n, n)(x, m, n) = \begin{cases} (e\theta^n \cdot x\theta^{n-m}, n, n + n - m) & \text{if } n \geq m \\ (e\theta^m \cdot x, m, n) & \text{if } n \leq m \end{cases}$$

$$(e\theta^n, n, n)(y, p, q) = \begin{cases} (e\theta^n \cdot y\theta^{n-p}, n, n + q - p) & \text{if } n \geq p \\ (e\theta^m \cdot y, p, q) & \text{if } n \leq p \end{cases}$$

From  $(e\theta^n, n, n)(x, m, n)$  and  $(e\theta^n, n, n)(y, p, q)$ , it follows that

$$n - (n + n - m) = m - n \text{ and } n - (n + q - p) = p - q.$$

It follows from definition that  $m - n = p - q$ .

For the first outer part of  $(e\theta^n, n, n)(x, m, n)$  and  $(e\theta^n, n, n)(y, p, q)$ , we have

$$\begin{aligned} e\theta^n \cdot x\theta^{n-m} &= e\theta^n \cdot y\theta^{n-p} \quad (\text{since from definition, } x\theta^{n-m} = y\theta^{n-p}) \\ e\theta^m \cdot x &= e\theta^p \cdot y \quad (\text{since from definition, } e\theta^{m-n} \cdot x = e\theta^{p-n} \cdot y). \end{aligned}$$

Thus  $(x, m, n) \mu_L (y, p, q)$  implies  $(e, 0, 0)(x, m, n) \mu_L (e, 0, 0)(y, p, q)$ .

To show that  $\mu \subseteq \mathcal{L}^*$ , we now consider the elements  $(x, m, n), (y, p, q) \in GBR^*(M, \theta)$  such that  $(x, m, n) \mu_L (y, p, q)$ . But  $(x, m, n)^* = (y, p, q)^*$  which implies that  $(x, m, n) \mathcal{L}^* (y, p, q)$ .

Now let  $\rho$  be a congruence on  $GBR^*(M, \theta)$  such that  $\rho \subseteq \mathcal{L}^*$ . If  $(x, m, n) \rho (y, p, q)$ , then for any  $(e, n, n) \in S$ ,  $(e, n, n)(x, m, n) \rho (e, n, n)(y, p, q)$  so that  $(e, n, n)(x, m, n) \mathcal{L}^* (e, n, n)(y, p, q)$ , that is  $(x, m, n) \mu_L (y, p, q)$  and whence  $\rho \subseteq \mu_L$ .

**Proposition 5.2.** Let  $S = GBR^*(M, \theta)$ . Then  $\mu_R$  is the maximum congruence on  $S$  contained in  $\mathcal{R}^*$ .

**Proof.** The proof is similar to the proof of Proposition 3.1.

An immediate consequence of Proposition 3.1 and Proposition 3.2 is the following

**Theorem 5.3.** Let  $S$  be a \*-bisimple type A I-semigroup. Then  $\mu$  is the maximum congruence on  $S$  contained in  $\mathcal{H}^*$ .

### Conflict of Interests

The authors declare that there is no conflict of interests.

### Acknowledgements

This work was partially supported by Joy Ndubuisi of Aba North Local Government, Department of Administration. The referee's comments have contributed substantially towards improving the presentation of the paper.

## REFERENCES

- [1] Asibong-Ibe, U.  $\ast$ -Bisimple type  $A$   $\omega$ -semigroups-1, *Semigroup Forum* 31 (1985), 99-117.
- [2] El-Qallali, A and Fountain, J.B. Idempotent-connected abundant semigroups. *Proc. Roy. Soc. Edinburgh Math. Sect. A* (1981), 79-90.
- [3] Fountain, J.B. A class of right PP monoids, *Q. J. Math. Oxford* 2 (28) (1974), 28-44.
- [4] Fountain, J.B. Adequate semigroups. *Proc. Edinburgh Math. Soc.* 22 (1979), 113-125.
- [5] Fountain, J.B. Abundant semigroups, *Proc. London. Math. Soc.* 3 (1) (1982), 103-129.
- [6] Howie, J.M. *Fundamentals of Semigroup Theory*, Oxford University Press, Inc. USA, 1995.
- [7] Makanjuola, S. O. Congruences on  $\ast$ -bisimple type  $A$   $\omega$ -semigroup. *J. Nigerian Math. Soc.* 11 (2) (1992), 21-28.
- [8] Piochi, B. Congruences on Bruck-Reilly extension of monoids. *Semigroup Forum*, 50 (1995), 179-191.
- [9] Shang, Y. and Wang, L.  $\ast$ -Bisimple type  $A$   $I$ -semigroups, *Southeast Asian Bull. Math.* 36 (2012), 535-545.
- [10] Warne, R.J.  $I$ -Bisimple semigroups. *Trans. Amer. Math. Soc.* 130 (1968), 367-386.